

# Nonradial normalized solutions for nonlinear scalar field equations

Louis Jeanjean\*

*Laboratoire de Mathématiques (CNRS UMR 6623)*

*Université de Bourgogne Franche-Comté*

*Besançon 25030, France*

*e-mail: louis.jeanjean@univ-fcomte.fr*

Sheng-Sen Lu†

*Center for Applied Mathematics, Tianjin University*

*Tianjin 300072, PR China*

*e-mail: sslu@tju.edu.cn*

**Abstract.** We study the following nonlinear scalar field equation

$$\begin{cases} -\Delta u = f(u) - \mu u & \text{in } \mathbb{R}^N, \\ \|u\|_{L^2(\mathbb{R}^N)}^2 = m, \\ u \in H^1(\mathbb{R}^N). \end{cases}$$

Here  $f \in C(\mathbb{R}, \mathbb{R})$ ,  $m > 0$  is a given constant and  $\mu \in \mathbb{R}$  is a Lagrange multiplier. In a mass subcritical case but under general assumptions on the nonlinearity  $f$ , we show the existence of one nonradial solution for any  $N \geq 4$ , and obtain multiple (sometimes infinitely many) nonradial solutions when  $N = 4$  or  $N \geq 6$ . In particular, all these solutions are sign-changing.

**2010 MSC:** 35J60, 58E05

**Key words:** Nonlinear scalar field equations,  $L^2$ -subcritical case, Nonradial solutions, Sign-changing solutions.

## Contents

1	<a href="#">Introduction</a>	2
2	<a href="#">A minimax theorem</a>	5
3	<a href="#">Preliminary results</a>	8
4	<a href="#">Proofs of the main results</a>	13
	4.1 <a href="#">Proof of Theorem 1.2</a> .....	13
	4.2 <a href="#">Proof of Theorem 1.1</a> .....	16
5	<a href="#">Another proof for Theorem B</a>	20
	<a href="#">References</a>	22

---

\*For L. Jeanjean this work has been carried out in the framework of the project NONLOCAL (ANR-14-CE25-0013) funded by the French National Research Agency (ANR).

†S.-S. Lu acknowledges the support of the NSF of China (NSFC-11771324), of the China Scholarship Council (CSC-201706250149) and the hospitality of the Laboratoire de Mathématiques (CNRS UMR 6623), Université de Bourgogne Franche-Comté.

# 1 Introduction

In this paper we consider the nonlinear scalar field equation with an  $L^2$  constraint:

$$\begin{cases} -\Delta u = f(u) - \mu u & \text{in } \mathbb{R}^N, \\ \|u\|_{L^2(\mathbb{R}^N)}^2 = m, \\ u \in H^1(\mathbb{R}^N). \end{cases} \quad (P_m)$$

Here  $N \geq 1$ ,  $f \in C(\mathbb{R}, \mathbb{R})$ ,  $m > 0$  is a given constant and  $\mu \in \mathbb{R}$  is a Lagrange multiplier.

The main feature of  $(P_m)$  is that the desired solutions have an a priori prescribed  $L^2$ -norm. Solutions of this type are often referred to as *normalized solutions* and they are interesting to be looked at from a physical point of view. Actually, one of the motivations to the search of solutions to equations of the type

$$-\Delta u = f(u) - \mu u, \quad u \in H^1(\mathbb{R}^N) \quad (1.1)$$

is that they correspond to stationary solutions of some nonlinear Klein-Gordon or Schrödinger type equations. We refer to the classical paper [5] in that direction. For these equations, finding solutions with a prescribed  $L^2$ -norm is relevant since this quantity is preserved along the time evolution.

Let  $I : H^1(\mathbb{R}^N) \rightarrow \mathbb{R}$  be the functional defined (at least formally) by

$$I(u) := \frac{1}{2} \int_{\mathbb{R}^N} |\nabla u|^2 dx - \int_{\mathbb{R}^N} F(u) dx,$$

where  $F(t) := \int_0^t f(\tau) d\tau$  for  $t \in \mathbb{R}$ . Clearly, solutions of  $(P_m)$  can be characterized as critical points of  $I$  submitted to the constraint

$$S_m := \left\{ u \in H^1(\mathbb{R}^N) \mid \|u\|_{L^2(\mathbb{R}^N)}^2 = m \right\}.$$

For future reference, for any  $u \in S_m$ , the value  $I(u)$  is called the *energy* of  $u$ ; if a solution of  $(P_m)$  minimizes  $I$  among all the solutions, we name it a *ground state*.

It is well-known that the study of  $(P_m)$  and the type of results one can expect, do depend on the behavior of the nonlinearity  $f$  at infinity. In particular, this behavior determines whether  $I$  is bounded from below on  $S_m$ . One speaks of a mass subcritical case if  $I$  is bounded from below on  $S_m$  for any  $m > 0$ , and of a mass supercritical case if  $I$  is unbounded from below on  $S_m$  for any  $m > 0$ . One also refers to a mass critical case when the boundedness from below does depend on the value  $m > 0$ . In this paper we focus on mass subcritical cases and we refer to the papers [1, 2, 11] for results in the mass supercritical cases.

The study of  $(P_m)$ , in the mass subcritical case and under appropriate conditions on  $f$ , started with the works of C.A. Stuart in the eighties [22, 23]. Concerning the existence of a ground state the more complete result is due to Shibata [21]. To be precise the following conditions were introduced on  $f$  in that paper:

$$(f1) \quad f \in C(\mathbb{R}, \mathbb{R}),$$

$$(f2) \quad \lim_{t \rightarrow 0} f(t)/t = 0,$$

$$(f3) \quad \lim_{t \rightarrow \infty} f(t)/|t|^{1+4/N} = 0,$$

$$(f4) \quad \text{There exists } \zeta > 0 \text{ such that } F(\zeta) > 0.$$

Since under (f1) – (f3) the functional  $I$  is bounded from below on  $S_m$ , one can define the global infimum

$$E_m := \inf_{u \in S_m} I(u) > -\infty$$

and expect the existence of a global minimizer and thus of a ground state to  $(P_m)$ . Indeed, Shibata proved the following existence result.

**Theorem A (Shibata [21])** *Assume that  $N \geq 1$  and  $f$  satisfies (f1) – (f4). Then there exists a uniquely determined number  $m_* \in [0, \infty)$  such that*

$$E_m = 0 \quad \text{if } 0 < m \leq m_*, \quad E_m < 0 \quad \text{when } m > m_*.$$

Moreover,

(i) *when  $m > m_*$ , the infimum  $E_m$  is achieved and thus  $(P_m)$  has a ground state;*

(ii)  *$m_* = 0$  if in addition*

$$\lim_{t \rightarrow 0} \frac{F(t)}{|t|^{2+\frac{4}{N}}} = +\infty; \tag{1.2}$$

(iii) *if  $0 < m < m_*$ , then the infimum  $E_m = 0$  is not achieved.*

Note also that [21] contains an orbital stability result for the set of minimizers when  $m > m_*$ . The derivation of such result relies on proving that any minimizing sequence to  $E_m$  is compact up to translations in  $\mathbb{R}^N$ , see [21] for more details.

Concerning the existence of more than one solution, the particular case  $f(u) = |u|^\sigma u$  with  $0 < \sigma < 4/N$  and  $N \geq 2$  was considered in [10] where infinitely many solutions (with negative energies) were obtained. For the most general result we refer to the recent paper [9] by Hirata and Tanaka. Still under the conditions (f1) – (f4), Hirata and Tanaka investigated  $(P_m)$  when  $N \geq 2$  and showed the existence of *multiple radial solutions* assuming in addition the oddness of  $f$ :

(f5)  $f(-t) = -f(t)$  for all  $t \in \mathbb{R}$ .

More precisely, applying a version of *symmetric mountain pass argument* to  $I(\lambda, u) : \mathbb{R} \times H_r^1(\mathbb{R}^N) \rightarrow \mathbb{R}$ , a Lagrange formulation of  $(P_m)$  defined as

$$I(\lambda, u) = \frac{1}{2} \int_{\mathbb{R}^N} |\nabla u|^2 dx - \int_{\mathbb{R}^N} F(u) dx + \frac{1}{2} e^\lambda \left( \int_{\mathbb{R}^N} u^2 dx - m \right),$$

they established Theorem B stated below.

**Theorem B (Hirata-Tanaka [9])** *Assume that  $N \geq 2$  and  $f$  satisfies (f5) in addition to (f1) – (f4). Then the following statements hold.*

(i) *For each  $k \in \mathbb{N}$  there exists  $\bar{m}_k \in [0, \infty)$  such that, when  $m > \bar{m}_k$ ,  $(P_m)$  has at least  $k$  radial solutions (with negative energies).*

(ii) *Assume in addition (1.2), then  $(P_m)$  has infinitely many radial solutions  $\{v_n\}_{n=1}^\infty$  for all  $m > 0$ . In particular,  $I(v_n) < 0$  for each  $n \in \mathbb{N}$  and  $I(v_n) \rightarrow 0$  as  $n \rightarrow \infty$ .*

We remark that just assuming (f1) – (f4), Hirata and Tanaka established the existence of one radial solution via a *mountain pass argument* applied to  $I(\lambda, u)$ . As a consequence they derived a minimax characterization of the infimum  $E_m$ , see [9, Theorem 0.1] for more details.

In this paper, we make further progress in the understanding of the set of solutions to  $(P_m)$ . Roughly speaking, when  $N \geq 4$  and under the conditions  $(f1) - (f5)$ , we derive existence and multiplicity results for *nonradial solutions* to  $(P_m)$ .

To state our results, we introduce some notations. Assume that  $N \geq 4$  and  $2 \leq M \leq N/2$ . Let us fix  $\tau \in \mathcal{O}(N)$  such that  $\tau(x_1, x_2, x_3) = (x_2, x_1, x_3)$  for  $x_1, x_2 \in \mathbb{R}^M$  and  $x_3 \in \mathbb{R}^{N-2M}$ , where  $x = (x_1, x_2, x_3) \in \mathbb{R}^N = \mathbb{R}^M \times \mathbb{R}^M \times \mathbb{R}^{N-2M}$ . We define

$$X_\tau := \{u \in H^1(\mathbb{R}^N) \mid u(\tau x) = -u(x) \text{ for all } x \in \mathbb{R}^N\}.$$

It is clear that  $X_\tau$  does not contain nontrivial radial functions. Let  $H_{\mathcal{O}_1}^1(\mathbb{R}^N)$  denote the subspace of invariant functions with respect to  $\mathcal{O}_1$ , where  $\mathcal{O}_1 := \mathcal{O}(M) \times \mathcal{O}(M) \times \text{id} \subset \mathcal{O}(N)$  acts isometrically on  $H^1(\mathbb{R}^N)$ . We also consider  $\mathcal{O}_2 := \mathcal{O}(M) \times \mathcal{O}(M) \times \mathcal{O}(N-2M) \subset \mathcal{O}(N)$  acting isometrically on  $H^1(\mathbb{R}^N)$  with the subspace of invariant functions denoted by  $H_{\mathcal{O}_2}^1(\mathbb{R}^N)$ . Here we agree that the components corresponding to  $N-2M$  do not exist when  $N=2M$ . Clearly,  $H_{\mathcal{O}_2}^1(\mathbb{R}^N)$  is in general a subspace of  $H_{\mathcal{O}_1}^1(\mathbb{R}^N)$ , but  $H_{\mathcal{O}_2}^1(\mathbb{R}^N) = H_{\mathcal{O}_1}^1(\mathbb{R}^N)$  when  $N=2M$ .

For notational convenience, we set  $X_1 := H_{\mathcal{O}_1}^1(\mathbb{R}^N) \cap X_\tau$  and  $X_2 := H_{\mathcal{O}_2}^1(\mathbb{R}^N) \cap X_\tau$ . Our first main result concerns the existence of one nonradial solution to  $(P_m)$ .

**Theorem 1.1** *Assume that  $N \geq 4$  and  $f$  satisfies  $(f1) - (f5)$ . Define*

$$E_{1,m} := \inf_{u \in S_m \cap X_1} I(u).$$

*Then  $E_{1,m} > -\infty$  and the mapping  $m \mapsto E_{1,m}$  is nonincreasing and continuous. Moreover*

*(i) there exists a uniquely determined number  $m^* \in [0, \infty)$  such that*

$$E_{1,m} = 0 \quad \text{if } 0 < m \leq m^*, \quad E_{1,m} < 0 \quad \text{when } m > m^*;$$

*(ii) when  $m > m^*$ , the infimum  $E_{1,m}$  is achieved and thus  $(P_m)$  has one nonradial solution  $w \in X_1$  such that  $I(w) = E_{1,m}$ ;*

*(iii) when  $0 < m < m^*$ ,  $E_{1,m}$  is not achieved;*

*(iv)  $m^* = 0$  if in addition (1.2) holds.*

Our second main result concerns the multiplicity of nonradial solutions to  $(P_m)$ . Let  $\Sigma(S_m \cap X_2)$  be the family of closed symmetric subsets of  $S_m \cap X_2$  and denote by  $\mathcal{G}(A)$  the genus of  $A \in \Sigma(S_m \cap X_2)$ .

**Theorem 1.2** *Assume that  $N \geq 4$ ,  $N-2M \neq 1$  and  $f$  satisfies  $(f1) - (f5)$ . Define the minimax values*

$$E_{m,k} := \inf_{A \in \Gamma_{m,k}} \sup_{u \in A} I(u),$$

*where  $\Gamma_{m,k} := \{A \in \Sigma(S_m \cap X_2) \mid \mathcal{G}(A) \geq k\}$ . Then the following statements hold.*

*(i)  $-\infty < E_{m,k} \leq E_{m,k+1} \leq 0$  for all  $m > 0$  and  $k \in \mathbb{N}$ .*

*(ii) For any  $k \in \mathbb{N}$ , the mapping  $m \mapsto E_{m,k}$  is nonincreasing and continuous.*

*(iii) For each  $k \in \mathbb{N}$ , there exists a uniquely determined  $m_k \in [0, \infty)$  such that*

$$E_{m,k} = 0 \quad \text{if } 0 < m \leq m_k, \quad E_{m,k} < 0 \quad \text{when } m > m_k.$$

*When  $m > m_k$ ,  $(P_m)$  has  $k$  distinct nonradial solutions belonging to  $X_2$  and associated to the levels  $E_{m,j}$  ( $j = 1, 2, \dots, k$ ).*

(iv) Assume in addition (1.2), then  $m_k = 0$  for any  $k \in \mathbb{N}$  and thus  $(P_m)$  has infinitely many nonradial solutions  $\{w_k\}_{k=1}^\infty \subset X_2$  for all  $m > 0$ . In particular,  $I(w_k) = E_{m,k} < 0$  for each  $k \in \mathbb{N}$  and  $I(w_k) \rightarrow 0$  as  $k \rightarrow \infty$ .

The question of the existence of nonradial solutions to the free equation (1.1) was raised in [5, Section 10.8] and remained open for a long time. Partial results, namely for specific nonlinearities  $f$ , were first obtained by Bartsch and Willem [3]. These authors worked in dimension  $N = 4$  and  $N \geq 6$  assuming an Ambrosetti-Rabinowitz type condition. Actually the idea of considering subspaces of  $H^1(\mathbb{R}^N)$  as  $H_{\mathcal{O}_2}^1(\mathbb{R}^N)$  originates from [3]. Note also the work [16] in which the problem is solved when  $N = 5$  by introducing the  $\mathcal{O}_1$  action on  $H^1(\mathbb{R}^N)$ . Finally it was only very recently that, under general assumptions on  $f$ , a positive answer to the existence and multiplicity of nonradial solutions was given [17]. Note also that in [12] the authors gave an alternative proof of the results of [17] with more elementary arguments. All these results however consider the equation (1.1) without prescribing the  $L^2$ -norm of the solutions.

The present paper is, up to our knowledge, the first to consider the existence of nonradial normalized solutions. We also observe that the nonradial solutions given by Theorems 1.1 and 1.2 change signs. In sharp contrast to the unconstrained case (1.1) where numerous results have been established, see for example [4, 15, 18], the existence of *sign-changing solutions* had not been studied yet for  $L^2$ -constrained problems.

Let us now give some ideas of the proofs of Theorems 1.1 and 1.2. To prove the multiplicity result stated in Theorem 1.2, we work in the space  $X_2 := H_{\mathcal{O}_2}^1(\mathbb{R}^N) \cap X_\tau$  and make use of classical minimax arguments (see Theorem 2.1 below). Since  $N \geq 4$  and  $N - 2M \neq 1$ , we can benefit from the compact inclusion  $X_2 \hookrightarrow L^p(\mathbb{R}^N)$  for all  $2 < p < 2N/(N - 2)$ . Indeed this result, which is due to P. L. Lions [14], allows to show that  $I_{|S_m \cap X_2}$  satisfies the Palais-Smale condition at any level  $c < 0$ , see Lemma 3.5. Another key point is to verify that the minimax levels  $E_{m,k}$  are indeed negative for some (or any)  $m > 0$  and  $k \in \mathbb{N}$ . Relying on the construction of some special mappings done in [12] (see also [17]), we manage to do this in Lemmas 3.4 and 4.1. Note also that to derive the existence of  $m_k$  in Theorem 1.2 (iii), we need Theorem 1.2 (ii) which is proved in Lemma 4.1.

For the proof of Theorem 1.1, we work in the space  $X_1 := H_{\mathcal{O}_1}^1(\mathbb{R}^N) \cap X_\tau$ . In the case where  $N - 2M = 0$ , we have  $X_1 = X_2$  (with  $N - 2M \neq 1$ ). Since then  $E_{1,m} = E_{m,1}$  and  $m^* = m_1$ , the existence of a minimizer for  $E_{1,m}$  when  $m > m^*$  follows directly from Theorem 1.2. When  $N - 2M \neq 0$ , the inclusion  $X_1 \hookrightarrow L^p(\mathbb{R}^N)$  is not compact for any  $2 < p < 2N/(N - 2)$  and the Palais-Smale condition does not hold any more. To derive our existence result in this case, using concentration compactness type arguments, we study carefully the behaviour of the minimizing sequences of  $E_{1,m}$ . In that direction we make use of some arguments from [21].

The paper is organized as follows. In Section 2 we present the version of the minimax theorem that will be used in the proof of Theorem 1.2. Section 3 establishes some key technical points to be used in the proofs of the main results. In Section 4 we prove Theorems 1.1 and 1.2. Finally, with the approach used to prove Theorem 1.2, we give in Section 5 an alternative proof for Theorem B, the multiplicity result due to Hirata and Tanaka [9].

## 2 A minimax theorem

In this section, we present a minimax theorem for a class of constrained even functionals. Let us point out that closely related results do exist in the literature, see in particular e.g., [6, Section 8], [20], [24] and [25, Chapter 5]. The present version is well suited to deal with the nonlinear scalar field equations considered in this paper.

To formulate the minimax theorem, we need some notations. Let  $\mathcal{E}$  be a real Banach space with norm  $\|\cdot\|_{\mathcal{E}}$  and  $\mathcal{H}$  be a real Hilbert space with inner product  $(\cdot, \cdot)_{\mathcal{H}}$ . We identify  $\mathcal{H}$  with its dual space and assume that  $\mathcal{E}$  is embedded continuously in  $\mathcal{H}$ . For any  $m > 0$ , define the manifold

$$\mathcal{M} := \{u \in \mathcal{E} \mid (u, u)_{\mathcal{H}} = m\},$$

which is endowed with the topology inherited from  $\mathcal{E}$ . Clearly, the tangent space of  $\mathcal{M}$  at a point  $u \in \mathcal{M}$  is defined by

$$T_u\mathcal{M} := \{v \in \mathcal{E} \mid (u, v)_{\mathcal{H}} = 0\}.$$

Let  $I \in C^1(\mathcal{E}, \mathbb{R})$ , then  $I|_{\mathcal{M}}$  is a functional of class  $C^1$  on  $\mathcal{M}$ . The norm of the derivative of  $I|_{\mathcal{M}}$  at any point  $u \in \mathcal{M}$  is defined by

$$\|I'_{|_{\mathcal{M}}}(u)\| := \sup_{\|v\|_{\mathcal{E}} \leq 1, v \in T_u\mathcal{M}} |\langle I'(u), v \rangle|.$$

A point  $u \in \mathcal{M}$  is said to be a critical point of  $I|_{\mathcal{M}}$  if  $I'_{|_{\mathcal{M}}}(u) = 0$  (or, equivalently,  $\|I'_{|_{\mathcal{M}}}(u)\| = 0$ ). A number  $c \in \mathbb{R}$  is called a critical value of  $I|_{\mathcal{M}}$  if  $I|_{\mathcal{M}}$  has a critical point  $u \in \mathcal{M}$  such that  $c = I(u)$ . We say that  $I|_{\mathcal{M}}$  satisfies the Palais-Smale condition at a level  $c \in \mathbb{R}$ ,  $(PS)_c$  for short, if any sequence  $\{u_n\} \subset \mathcal{M}$  with  $I(u_n) \rightarrow c$  and  $\|I'_{|_{\mathcal{M}}}(u_n)\| \rightarrow 0$  contains a convergent subsequence.

Noting that  $\mathcal{M}$  is symmetric with respect to  $0 \in \mathcal{E}$  and  $0 \notin \mathcal{M}$ , we introduce the notation of the genus. Let  $\Sigma(\mathcal{M})$  be the family of closed symmetric subsets of  $\mathcal{M}$ . For any nonempty set  $A \in \Sigma(\mathcal{M})$ , the genus  $\mathcal{G}(A)$  of  $A$  is defined as the least integer  $k \geq 1$  for which there exists an odd continuous mapping  $\varphi : A \rightarrow \mathbb{R}^k \setminus \{0\}$ . We set  $\mathcal{G}(A) = \infty$  if such an integer does not exist, and set  $\mathcal{G}(A) = 0$  if  $A = \emptyset$ . For each  $k \in \mathbb{N}$ , let  $\Gamma_k := \{A \in \Sigma(\mathcal{M}) \mid \mathcal{G}(A) \geq k\}$ .

We now state the minimax theorem and then give a detailed proof for completeness.

**Theorem 2.1 (Minimax theorem)** *Let  $I : \mathcal{E} \rightarrow \mathbb{R}$  be an even functional of class  $C^1$ . Assume that  $I|_{\mathcal{M}}$  is bounded from below on  $\mathcal{M}$  and satisfies the condition  $(PS)_c$  for all  $c < 0$ , and that  $\Gamma_k \neq \emptyset$  for each  $k \in \mathbb{N}$ . Then a sequence of minimax values  $-\infty < c_1 \leq c_2 \leq \dots \leq c_k \leq \dots$  can be defined as follows:*

$$c_k := \inf_{A \in \Gamma_k} \sup_{u \in A} I(u), \quad k \geq 1,$$

and the following statements hold.

(i)  $c_k$  is a critical value of  $I|_{\mathcal{M}}$  provided  $c_k < 0$ .

(ii) Denote by  $K^c$  the set of critical points of  $I|_{\mathcal{M}}$  at a level  $c \in \mathbb{R}$ . If

$$c_k = c_{k+1} = \dots = c_{k+l-1} =: c < 0 \quad \text{for some } k, l \geq 1,$$

then  $\mathcal{G}(K^c) \geq l$ . In particular,  $I|_{\mathcal{M}}$  has infinitely many critical points at the level  $c$  if  $l \geq 2$ .

(iii) If  $c_k < 0$  for all  $k \geq 1$ , then  $c_k \rightarrow 0^-$  as  $k \rightarrow \infty$ .

To prove Theorem 2.1, we shall need some basic properties of the genus. For  $A \subset \mathcal{M}$  and  $\delta > 0$ , denote by  $A_\delta$  the uniform  $\delta$ -neighborhood of  $A$  in  $\mathcal{M}$ , that is,

$$A_\delta := \{u \in \mathcal{M} \mid \inf_{v \in A} \|u - v\|_{\mathcal{E}} \leq \delta\}.$$

Since  $\mathcal{M}$  is a closed symmetric subset of  $\mathcal{E}$ , repeating the arguments in [20, Section 7], one can get Proposition 2.2 below which is sufficient for our use.

**Proposition 2.2** *Let  $A, B \in \Sigma(\mathcal{M})$ . Then the following statements hold.*

- (i) *If  $\mathcal{G}(A) \geq 2$ , then  $A$  contains infinitely many distinct points.*
- (ii)  *$\mathcal{G}(\overline{A \setminus B}) \geq \mathcal{G}(A) - \mathcal{G}(B)$  if  $\mathcal{G}(B) < \infty$ .*
- (iii) *If there exists an odd continuous mapping  $\psi : \mathbb{S}^{k-1} \rightarrow A$ , then  $\mathcal{G}(A) \geq k$ .*
- (iv) *If  $A$  is compact, then  $\mathcal{G}(A) < \infty$  and there exists  $\delta > 0$  such that  $A_\delta \in \Sigma(\mathcal{M})$  and  $\mathcal{G}(A_\delta) = \mathcal{G}(A)$ .*

We shall also need the following quantitative deformation lemma whose proof is similar to that of [25, Lemma 2.3]. For  $c < d$ , set  $I_{|\mathcal{M}}^c := \{u \in \mathcal{M} \mid I(u) \leq c\}$  and  $I_{|\mathcal{M}}^{-1}([c, d]) := \{u \in \mathcal{M} \mid c \leq I(u) \leq d\}$ .

**Lemma 2.3** *Assume  $I_{|\mathcal{M}} \in C^1(\mathcal{M}, \mathbb{R})$ . Let  $S \subset \mathcal{M}$ ,  $c \in \mathbb{R}$ ,  $\varepsilon > 0$  and  $\delta > 0$  such that*

$$\|I'_{|\mathcal{M}}(u)\| \geq \frac{8\varepsilon}{\delta} \quad \text{for all } u \in I_{|\mathcal{M}}^{-1}([c - 2\varepsilon, c + 2\varepsilon]) \cap S_{2\delta}. \quad (2.1)$$

*Then there exists a mapping  $\eta \in C([0, 1] \times \mathcal{M}, \mathcal{M})$  such that*

- (i)  *$\eta(t, u) = u$  if  $t = 0$  or if  $u \notin I_{|\mathcal{M}}^{-1}([c - 2\varepsilon, c + 2\varepsilon]) \cap S_{2\delta}$ ,*
- (ii)  *$\eta(1, I_{|\mathcal{M}}^{c+\varepsilon} \cap S) \subset I_{|\mathcal{M}}^{c-\varepsilon}$ ,*
- (iii)  *$I(\eta(t, u))$  is nonincreasing in  $t \in [0, 1]$  for any  $u \in \mathcal{M}$ ,*
- (iv)  *$\eta(t, u)$  is odd in  $u \in \mathcal{M}$  for any  $t \in [0, 1]$  if  $I_{|\mathcal{M}}$  is even.*

With Proposition 2.2 and Lemma 2.3 in hand, we can now prove Theorem 2.1.

**Proof of Theorem 2.1.** Item (i) is a special case of Item (ii) when  $l = 1$ , so we go straight to the proof of Item (ii). Obviously,  $K^c \in \Sigma(\mathcal{M})$  and  $K^c$  is compact by the condition  $(PS)_c$ . If  $\mathcal{G}(K^c) \leq l - 1$ , by Proposition 2.2 (iv), there exists  $\delta > 0$  such that

$$\mathcal{G}(K_{3\delta}^c) = \mathcal{G}(K^c) \leq l - 1.$$

We remark here that  $K_{3\delta}^c = \emptyset$  if  $K^c = \emptyset$ . Let  $S := \overline{\mathcal{M} \setminus K_{3\delta}^c} \subset \mathcal{M}$ . Clearly, there exists  $\varepsilon > 0$  small enough such that (2.1) holds (if not, one will get a Palais-Smale sequence  $\{u_n\} \subset S_{2\delta}$  of  $I_{|\mathcal{M}}$  at the level  $c < 0$  and thus an element  $v \in S_{2\delta} \cap K^c$  by the condition  $(PS)_c$ , which leads a contradiction since  $S_{2\delta} \cap K^c = \emptyset$ ). Therefore, Lemma 2.3 yields a mapping  $\eta \in C([0, 1] \times \mathcal{M}, \mathcal{M})$  such that

$$\eta(1, I_{|\mathcal{M}}^{c+\varepsilon} \cap S) \subset I_{|\mathcal{M}}^{c-\varepsilon} \quad \text{and} \quad \eta(t, \cdot) \text{ is odd for all } t \in [0, 1].$$

Choose  $A \in \Gamma_{k+l-1}$  such that  $\sup_{u \in A} I(u) \leq c + \varepsilon$ . It is clear that  $\overline{A \setminus K_{3\delta}^c} \subset I_{|\mathcal{M}}^{c+\varepsilon} \cap S$  and thus

$$\eta(1, \overline{A \setminus K_{3\delta}^c}) \subset \eta(1, I_{|\mathcal{M}}^{c+\varepsilon} \cap S) \subset I_{|\mathcal{M}}^{c-\varepsilon}. \quad (2.2)$$

On the other hand, since  $\mathcal{G}(\overline{A \setminus K_{3\delta}^c}) \geq \mathcal{G}(A) - \mathcal{G}(K_{3\delta}^c) \geq k$  by Proposition 2.2 (ii), we have  $\overline{A \setminus K_{3\delta}^c} \in \Gamma_k$  and then  $\eta(1, \overline{A \setminus K_{3\delta}^c}) \in \Gamma_k$ . Now, by the definition of  $c_k$  and (2.2), we get a contradiction:

$$c = c_k \leq \sup_{u \in \eta(1, \overline{A \setminus K_{3\delta}^c})} I(u) \leq c - \varepsilon.$$

Thus  $\mathcal{G}(K^c) \geq l$ . In view of Proposition 2.2 (i), we complete the proof of Item (ii).



To prove Item (iii), we assume by contradiction that there exists  $c < 0$  such that  $c_k \leq c$  for all  $k \geq 1$  and  $c_k \rightarrow c$  as  $k \rightarrow \infty$ . By the condition  $(PS)_c$ ,  $K^c$  is a (symmetric) compact set. Thus, by Proposition 2.2 (iv), there exists  $\delta > 0$  such that

$$\mathcal{G}(K_{3\delta}^c) = \mathcal{G}(K^c) =: q < \infty.$$

Let  $S := \overline{\mathcal{M} \setminus K_{3\delta}^c} \subset \mathcal{M}$ . Since (2.1) holds for small enough  $\varepsilon > 0$ , we know from Lemma 2.3 that a mapping  $\eta \in C([0, 1] \times \mathcal{M}, \mathcal{M})$  exists such that  $\eta(1, I_{|\mathcal{M}}^{c+\varepsilon} \cap S) \subset I_{|\mathcal{M}}^{c-\varepsilon}$  and  $\eta(t, \cdot)$  is odd for any  $t \in [0, 1]$ . Choose  $k \geq 1$  large enough such that  $c_k > c - \varepsilon$  and take  $A \in \Gamma_{k+q}$  such that  $\sup_{u \in A} I(u) \leq c_{k+q} + \varepsilon$ . Noting that  $c_{k+q} \leq c$ , we have  $\overline{A \setminus K_{3\delta}^c} \subset I_{|\mathcal{M}}^{c+\varepsilon} \cap S$  and thus

$$\eta(1, \overline{A \setminus K_{3\delta}^c}) \subset \eta(1, I_{|\mathcal{M}}^{c+\varepsilon} \cap S) \subset I_{|\mathcal{M}}^{c-\varepsilon}.$$

On the other hand, since  $\mathcal{G}(\overline{A \setminus K_{3\delta}^c}) \geq \mathcal{G}(A) - \mathcal{G}(K_{3\delta}^c) \geq k$ , we have  $\overline{A \setminus K_{3\delta}^c} \in \Gamma_k$  and then  $\eta(1, \overline{A \setminus K_{3\delta}^c}) \in \Gamma_k$ . We now reach a contradiction:

$$c_k \leq \sup_{u \in \eta(1, \overline{A \setminus K_{3\delta}^c})} I(u) \leq c - \varepsilon,$$

for we chosen  $k$  large enough such that  $c_k > c - \varepsilon$ . Thus  $c_k \rightarrow 0^-$  as  $k \rightarrow \infty$ .  $\square$

To end this section, we recall a characterization result in [6] which allows to check the condition  $(PS)_c$  in a convenient way.

**Lemma 2.4** ([6, Lemma 3]) *Assume that  $I : \mathcal{E} \rightarrow \mathbb{R}$  is of class  $C^1$ . Let  $\{u_n\}$  be a sequence in  $\mathcal{M}$  which is bounded in  $\mathcal{E}$ . Then the following are equivalent:*

(i)  $\|I'_{|\mathcal{M}}(u_n)\| \rightarrow 0$  as  $n \rightarrow \infty$ .

(ii)  $I'(u_n) - m^{-1} \langle I'(u_n), u_n \rangle u_n \rightarrow 0$  in  $\mathcal{E}^{-1}$  (the dual space of  $\mathcal{E}$ ) as  $n \rightarrow \infty$ .

Here the last  $u_n$  in (ii) is an element of  $\mathcal{E}^{-1}$  such that  $\langle u_n, v \rangle := (u_n, v)_{\mathcal{H}}$  for all  $v \in \mathcal{E}$ .

### 3 Preliminary results

In this section, we present some preliminary results. The first one is Lemma 3.1 below, which is a slightly modified version of [21, Lemma 2.2].

**Lemma 3.1** *Assume that  $N \geq 1$  and  $f$  satisfies (f1) – (f3). Then the following statements hold.*

(i) *Let  $\{u_n\}$  be a bounded sequence in  $H^1(\mathbb{R}^N)$ . We have*

$$\lim_{n \rightarrow \infty} \int_{\mathbb{R}^N} F(u_n) dx = 0$$

*if either  $\lim_{n \rightarrow \infty} \|u_n\|_{L^2(\mathbb{R}^N)} = 0$  or  $\lim_{n \rightarrow \infty} \|u_n\|_{L^{2+4/N}(\mathbb{R}^N)} = 0$ .*

(ii) *There exists  $C = C(f, N, m) > 0$  depending on  $f$ ,  $N$  and  $m > 0$  such that*

$$I(u) \geq \frac{1}{4} \int_{\mathbb{R}^N} |\nabla u|^2 dx - C(f, N, m) \tag{3.1}$$

*for all  $u \in H^1(\mathbb{R}^N)$  satisfying  $\|u\|_{L^2(\mathbb{R}^N)}^2 \leq m$ .*



**Lemma 3.2** *Assume that  $N \geq 2$ ,  $\{u_n\} \subset H^1(\mathbb{R}^N)$  is a bounded sequence, and  $u_n \rightarrow u$  almost everywhere in  $\mathbb{R}^N$  for some  $u \in H^1(\mathbb{R}^N)$ . Let  $F : \mathbb{R} \rightarrow \mathbb{R}$  be a function of class  $C^1$  with  $F(0) = 0$ . If*

(i) *when  $N = 2$ , for any  $\alpha > 0$ , there exists  $C_\alpha > 0$  such that*

$$|F'(t)| \leq C_\alpha \left[ |t| + \left( e^{\alpha t^2} - 1 \right) \right] \quad \text{for all } t \in \mathbb{R}.$$

(ii) *when  $N \geq 3$ , there exists  $C > 0$  such that  $|F'(t)| \leq C (|t| + |t|^{2^*-1})$  for all  $t \in \mathbb{R}$ ,*

then

$$\lim_{n \rightarrow \infty} \int_{\mathbb{R}^N} |F(u_n) - F(u_n - u) - F(u)| dx = 0. \quad (3.2)$$

**Proof.** We prove this lemma by applying the Brezis-Lieb Lemma ([7, Theorem 2]). Clearly,

$$|F(a+b) - F(a)| = \left| \int_0^1 \frac{d}{d\tau} F(a + \tau b) d\tau \right| = \left| \int_0^1 F'(a + \tau b) b d\tau \right| \quad \text{for all } a, b \in \mathbb{R}. \quad (3.3)$$

Let  $\varepsilon > 0$  be arbitrary. When  $N = 2$ , by (3.3), (i) and Young's inequality, one has

$$\begin{aligned} |F(a+b) - F(a)| &\leq C_\alpha \int_0^1 \left\{ |a + \tau b| + \left[ e^{\alpha(a+\tau b)^2} - 1 \right] \right\} |b| d\tau \\ &\leq C_\alpha \left[ |a| + |b| + \left( e^{4\alpha a^2} - 1 \right) + \left( e^{4\alpha b^2} - 1 \right) \right] |b| \\ &\leq C_\alpha \left[ \varepsilon a^2 + \varepsilon^{-1} b^2 + b^2 + \varepsilon \left( e^{4\alpha a^2} - 1 \right)^2 + \varepsilon^{-1} b^2 + \left( e^{4\alpha b^2} - 1 \right)^2 + b^2 \right] \\ &\leq \varepsilon C_\alpha \left[ a^2 + \left( e^{8\alpha a^2} - 1 \right) \right] + C_\alpha \left[ 2(1 + \varepsilon^{-1}) b^2 + \left( e^{8\alpha b^2} - 1 \right) \right] \\ &=: \varepsilon \varphi(a) + \psi_\varepsilon(b). \end{aligned}$$

In particular,  $|F(b)| \leq \psi_\varepsilon(b)$  for all  $b \in \mathbb{R}$ . Choose  $M \geq 1$  sufficiently large and  $\alpha > 0$  small enough such that

$$\|u_n\|_{H^1(\mathbb{R}^2)}, \|u_n - u\|_{H^1(\mathbb{R}^2)}, \|u\|_{H^1(\mathbb{R}^2)} \leq M$$

and

$$8\alpha \leq \frac{\beta}{M^2} \quad \text{for some } \beta \in (0, 4\pi).$$

By the Moser-Trudinger inequality, we know that  $\int_{\mathbb{R}^2} \varphi(u_n - u) dx$  is bounded uniformly in  $\varepsilon$  and  $n$ ,  $\int_{\mathbb{R}^2} \psi_\varepsilon(u) dx < \infty$  for any  $\varepsilon > 0$ , and  $F(u) \in L^1(\mathbb{R}^2)$ . In view of [7, Theorem 2], we obtain (3.2).

When  $N \geq 3$ , by (3.3), (ii) and Young's inequality, one has

$$\begin{aligned} |F(a+b) - F(a)| &\leq C \int_0^1 \left( |a + \tau b| + |a + \tau b|^{2^*-1} \right) |b| d\tau \\ &\leq C \left( |a| + 2^{2^*} |a|^{2^*-1} + |b| + 2^{2^*} |b|^{2^*-1} \right) |b| \\ &\leq \varepsilon C \left( a^2 + |2a|^{2^*} \right) + C \left[ (1 + \varepsilon^{-1}) b^2 + \left( 1 + \varepsilon^{1-2^*} \right) |2b|^{2^*} \right] \\ &=: \varepsilon \varphi(a) + \psi_\varepsilon(b). \end{aligned}$$

Applying Sobolev inequality and [7, Theorem 2], one can conclude easily that (3.2) holds.  $\square$

**Lemma 3.3** ([17, Corollary 3.2]) *Assume that  $N \geq 4$  and  $N - 2M \neq 0$ . Let  $\{u_n\}$  be a bounded sequence in  $H_{O_1}^1(\mathbb{R}^N)$  which satisfies*

$$\lim_{r \rightarrow \infty} \left( \lim_{n \rightarrow \infty} \sup_{y \in \{0\} \times \{0\} \times \mathbb{R}^{N-2M}} \int_{B(y,r)} |u_n|^2 dx \right) = 0. \quad (3.4)$$

Then  $u_n \rightarrow 0$  in  $L^p(\mathbb{R}^N)$  for any  $2 < p < 2^*$ .

**Proof.** We give here a complete proof for the reader's convenience. By [25, Lemma 1.21], the proof will be over if we can show that

$$\limsup_{n \rightarrow \infty} \sup_{y \in \mathbb{R}^N} \int_{B(y,1)} |u_n|^2 dx = 0. \quad (3.5)$$

We assume by contradiction that (3.5) does not hold. Thus, up to a subsequence, there exist  $\delta > 0$  and  $\{y_n\} \subset \mathbb{R}^N$  such that

$$\int_{B(y_n,1)} |u_n|^2 dx \geq \delta > 0 \quad \text{for } n \geq 1 \text{ large enough.} \quad (3.6)$$

Since  $\{u_n\}$  is bounded in  $L^2(\mathbb{R}^N)$  and invariant with respect to  $\mathcal{O}_1$ , in view of (3.6), we deduce that  $\{|(y_n^1, y_n^2)|\}$  must be bounded. Indeed, if  $|(y_n^1, y_n^2)| \rightarrow \infty$ , then one will derive the existence of an arbitrarily large number of disjoint unit balls in the family  $\{B(g^{-1}y_n, 1)\}_{g \in \mathcal{O}_1}$ . Thus, for sufficiently large  $r$ , we have

$$\int_{B((0,0,y_n^2),r)} |u_n|^2 dx \geq \int_{B(y_n,1)} |u_n|^2 dx \geq \delta > 0,$$

which contradicts (3.4). Therefore, (3.5) is satisfied and the desired conclusion follows.  $\square$

For any  $k \in \mathbb{N}$ , let  $\mathbb{S}^{k-1}$  be the unit sphere in  $\mathbb{R}^k$ , i.e.,

$$\mathbb{S}^{k-1} := \{x \in \mathbb{R}^k \mid |x| = 1\}.$$

Recall that  $X_2 := H_{\mathcal{O}_2}^1(\mathbb{R}^N) \cap X_\tau$ .

**Lemma 3.4** *Assume that  $N \geq 4$  and  $f$  satisfies (f1) – (f5). Then, for any  $m > 0$  and  $k \in \mathbb{N}$ , there exists an odd continuous mapping  $\gamma_{m,k} : \mathbb{S}^{k-1} \rightarrow S_m \cap X_2$ . Moreover, the following statements hold.*

(i) *For any  $k \in \mathbb{N}$ , there exists  $m(k) > 0$  large enough such that*

$$\sup_{\sigma \in \mathbb{S}^{k-1}} I(\gamma_{m,k}[\sigma]) < 0 \quad \text{for all } m > m(k). \quad (3.7)$$

(ii) *For any  $s > 0$ , define  $\gamma_{m,k}^s : \mathbb{S}^{k-1} \rightarrow S_m \cap X_2$  as follows:*

$$\gamma_{m,k}^s[\sigma](x) := s^{N/2} \gamma_{m,k}[\sigma](sx), \quad x \in \mathbb{R}^N \text{ and } \sigma \in \mathbb{S}^{k-1}.$$

*Then*

$$\limsup_{s \rightarrow 0^+} \left( \sup_{\sigma \in \mathbb{S}^{k-1}} I(\gamma_{m,k}^s[\sigma]) \right) \leq 0. \quad (3.8)$$

*If in addition (1.2) holds, then there exists  $s_* > 0$  small enough such that*

$$\sup_{\sigma \in \mathbb{S}^{k-1}} I(\gamma_{m,k}^s[\sigma]) < 0 \quad \text{for any } s \in (0, s_*). \quad (3.9)$$

**Proof.** Fix  $m > 0$  and  $k \in \mathbb{N}$ . Since  $f$  satisfies (f4), we know from [12, Eq. (4.3) and Lemma 4.3] (see also [17]) that there exists an odd continuous mapping  $\pi_k : \mathbb{S}^{k-1} \rightarrow X_2 \setminus \{0\}$  such that

$$\inf_{\sigma \in \mathbb{S}^{k-1}} \int_{\mathbb{R}^N} F(\pi_k[\sigma]) dx \geq 1 \quad \text{and} \quad \sup_{\sigma \in \mathbb{S}^{k-1}} \|\pi_k[\sigma]\|_{L^\infty(\mathbb{R}^N)} \leq 2\zeta.$$

Thus, we can define an odd continuous mapping  $\gamma_{m,k} : \mathbb{S}^{k-1} \rightarrow S_m \cap X_2$  as follows:

$$\gamma_{m,k}[\sigma](x) := \pi_k[\sigma] \left( m^{-1/N} \cdot \|\pi_k[\sigma]\|_{L^2(\mathbb{R}^N)}^{2/N} \cdot x \right), \quad x \in \mathbb{R}^N \text{ and } \sigma \in \mathbb{S}^{k-1}.$$

Clearly,

$$\sup_{\sigma \in \mathbb{S}^{k-1}} \|\gamma_{m,k}(\sigma)\|_{L^\infty(\mathbb{R}^N)} \leq 2\zeta.$$

We next show that this mapping  $\gamma_{m,k}$  satisfies Items (i) and (ii).

(i) Since  $\mathbb{S}^{k-1}$  is compact and  $0 \notin \pi_k[\mathbb{S}^{k-1}]$ , one can find  $\alpha_k, \beta_k, \beta'_k > 0$  independent of  $\sigma \in \mathbb{S}^{k-1}$  such that

$$\int_{\mathbb{R}^N} |\nabla \pi_k[\sigma]|^2 dx \leq \alpha_k \quad \text{and} \quad \beta_k \leq \|\pi_k[\sigma]\|_{L^2(\mathbb{R}^N)}^2 \leq \beta'_k.$$

Thus,

$$\begin{aligned} I(\gamma_{m,k}[\sigma]) &= \frac{1}{2} \int_{\mathbb{R}^N} |\nabla \gamma_{m,k}[\sigma]|^2 dx - \int_{\mathbb{R}^N} F(\gamma_{m,k}[\sigma]) dx \\ &= \frac{m^{\frac{N-2}{N}}}{2 \|\pi_k[\sigma]\|_{L^2(\mathbb{R}^N)}^{2(N-2)/N}} \int_{\mathbb{R}^N} |\nabla \pi_k[\sigma]|^2 dx - \frac{m}{\|\pi_k[\sigma]\|_{L^2(\mathbb{R}^N)}^2} \int_{\mathbb{R}^N} F(\pi_k[\sigma]) dx \\ &\leq \frac{1}{2} \alpha_k \beta_k^{(2-N)/N} \cdot m^{\frac{N-2}{N}} - (\beta'_k)^{-1} \cdot m := g_k(m). \end{aligned}$$

Clearly,  $g_k(m) < 0$  for sufficiently large  $m > 0$  and thus (3.7) holds.

(ii) We first prove (3.8). Let  $\varepsilon > 0$  be arbitrary. By (f2), there exists  $\delta > 0$  such that

$$|F(t)| \leq \varepsilon t^2 \quad \text{for all } |t| \leq \delta.$$

Noting that

$$\|\gamma_{m,k}^s[\sigma]\|_{L^\infty(\mathbb{R}^N)} \leq 2s^{N/2}\zeta \quad \text{for all } \sigma \in \mathbb{S}^{k-1}, \quad (3.10)$$

one can find  $s(\varepsilon) > 0$  small enough such that

$$\sup_{\sigma \in \mathbb{S}^{k-1}} \|\gamma_{m,k}^s[\sigma]\|_{L^\infty(\mathbb{R}^N)} \leq \delta \quad \text{for all } 0 < s < s(\varepsilon).$$

Therefore, for any  $\sigma \in \mathbb{S}^{k-1}$  and  $0 < s < s(\varepsilon)$ , we have

$$\begin{aligned} I(\gamma_{m,k}^s[\sigma]) &\leq \frac{1}{2} \int_{\mathbb{R}^N} |\nabla \gamma_{m,k}^s[\sigma]|^2 dx + \int_{\mathbb{R}^N} |F(\gamma_{m,k}^s[\sigma])| dx \\ &\leq \frac{1}{2} s^2 \int_{\mathbb{R}^N} |\nabla \gamma_{m,k}[\sigma]|^2 dx + \varepsilon \int_{\mathbb{R}^N} |\gamma_{m,k}[\sigma]|^2 dx \\ &= \frac{1}{2} s^2 \int_{\mathbb{R}^N} |\nabla \gamma_{m,k}[\sigma]|^2 dx + m\varepsilon. \end{aligned}$$

Since  $\mathbb{S}^{k-1}$  is compact, there exists  $C > 0$  (independent of  $\varepsilon > 0$  and  $s > 0$ ) such that

$$\sup_{\sigma \in \mathbb{S}^{k-1}} \int_{\mathbb{R}^N} |\nabla \gamma_{m,k}[\sigma]|^2 dx \leq C.$$

Thus, for any  $0 < s < \min\{s(\varepsilon), (2\varepsilon/C)^{1/2}\}$ , we obtain

$$\sup_{\sigma \in \mathbb{S}^{k-1}} I(\gamma_{m,k}^s[\sigma]) \leq \frac{1}{2} C s^2 + m\varepsilon \leq (m+1)\varepsilon.$$

Clearly, it follows that (3.8) holds.

We now assume (1.2) and prove (3.9). For

$$D := \sup_{\sigma \in \mathbb{S}^{k-1}} \int_{\mathbb{R}^N} |\nabla \gamma_{m,k}[\sigma]|^2 dx \Big/ \inf_{\sigma \in \mathbb{S}^{k-1}} \int_{\mathbb{R}^N} |\gamma_{m,k}[\sigma]|^{2+\frac{4}{N}} dx > 0,$$

by (1.2), there exists a  $\delta > 0$  such that

$$F(t) \geq D|t|^{2+\frac{4}{N}} \quad \text{for all } |t| < \delta.$$

Also, in view of (3.10), one can find  $s_* > 0$  small enough such that

$$\sup_{\sigma \in \mathbb{S}^{k-1}} \|\gamma_{m,k}^s[\sigma]\|_{L^\infty(\mathbb{R}^N)} \leq \delta \quad \text{for all } 0 < s < s_*.$$

Thus, for any  $\sigma \in \mathbb{S}^{k-1}$  and  $0 < s < s_*$ , we have

$$\begin{aligned} I(\gamma_{m,k}^s[\sigma]) &\leq \frac{1}{2} \int_{\mathbb{R}^N} |\nabla \gamma_{m,k}^s[\sigma]|^2 dx - D \int_{\mathbb{R}^N} |\gamma_{m,k}^s[\sigma]|^{2+\frac{4}{N}} dx \\ &= \frac{1}{2} s^2 \int_{\mathbb{R}^N} |\nabla \gamma_{m,k}[\sigma]|^2 dx - D s^2 \int_{\mathbb{R}^N} |\gamma_{m,k}[\sigma]|^{2+\frac{4}{N}} dx \\ &\leq -\frac{1}{2} s^2 \sup_{\sigma \in \mathbb{S}^{k-1}} \int_{\mathbb{R}^N} |\nabla \gamma_{m,k}[\sigma]|^2 dx < 0, \end{aligned}$$

which implies (3.9).  $\square$

**Lemma 3.5** *Assume that  $N \geq 4$ ,  $N - 2M \neq 1$ , and  $f$  satisfies (f1) – (f3) and (f5). Then  $I|_{S_m \cap X_2}$  satisfies the condition (PS) $_c$  for all  $c < 0$ .*

**Proof.** For given  $c < 0$ , let  $\{u_n\} \subset S_m \cap X_2$  be any sequence such that

$$I(u_n) \rightarrow c < 0 \tag{3.11}$$

and

$$I'|_{S_m \cap X_2}(u_n) \rightarrow 0. \tag{3.12}$$

By (3.11) and Lemma 3.1 (ii), we see that  $\{u_n\}$  is bounded in  $X_2$ . Thus, up to a subsequence, we may assume that  $u_n \rightharpoonup u$  in  $X_2$  and  $u_n \rightarrow u$  almost everywhere in  $\mathbb{R}^N$  for some  $u \in X_2$ . In addition, thanks to [25, Corollary 1.25],  $u_n \rightarrow u$  in  $L^p(\mathbb{R}^N)$  for any  $p \in (2, 2 + 4/N]$ . Also, we know from (3.12) and Lemma 2.4 that

$$-\Delta u_n + \mu_n u_n - f(u_n) \rightarrow 0 \quad \text{in } (X_2)^{-1}, \tag{3.13}$$

where

$$\mu_n := \frac{1}{m} \left( \int_{\mathbb{R}^N} f(u_n) u_n dx - \int_{\mathbb{R}^N} |\nabla u_n|^2 dx \right).$$

Since  $\{\mu_n\}$  is bounded by (f1) – (f3), we may assume that  $\mu_n \rightarrow \mu$  for some  $\mu \in \mathbb{R}$  and thus

$$-\Delta u + \mu u = f(u) \quad \text{in } (X_2)^{-1}. \tag{3.14}$$

To show that  $u_n \rightarrow u$  in  $X_2$ , the following two claims are needed.

**Claim 1.**

$$\lim_{n \rightarrow \infty} \int_{\mathbb{R}^N} f(u_n) u_n dx = \int_{\mathbb{R}^N} f(u) u dx. \tag{3.15}$$

Let  $v_n := u_n - u$  and  $q := 2 + 4/N$ . Clearly,

$$\int_{\mathbb{R}^N} [f(u_n) u_n - f(u) u] dx = \int_{\mathbb{R}^N} f(u_n) v_n dx + \int_{\mathbb{R}^N} [f(u_n) - f(u)] u dx.$$

Since  $u_n \rightharpoonup u$  in  $X_2$ , one can show in a standard way that  $\int_{\mathbb{R}^N} [f(u_n) - f(u)] u dx \rightarrow 0$ . We estimate the remaining term  $\int_{\mathbb{R}^N} f(u_n) v_n dx$ . For any  $\varepsilon > 0$ , there exists  $C_\varepsilon > 0$  such that

$$|f(t)| \leq \varepsilon |t| + C_\varepsilon |t|^{q-1} \quad \text{for all } t \in \mathbb{R}.$$

Therefore, by Hölder inequality, we have

$$\left| \int_{\mathbb{R}^N} f(u_n)v_n dx \right| \leq \varepsilon \|u_n\|_{L^2(\mathbb{R}^N)} \|v_n\|_{L^2(\mathbb{R}^N)} + C_\varepsilon \|u_n\|_{L^q(\mathbb{R}^N)}^{q-1} \|v_n\|_{L^q(\mathbb{R}^N)}.$$

Since  $\|v_n\|_{L^q(\mathbb{R}^N)} \rightarrow 0$  and  $\varepsilon > 0$  is arbitrary, we see that

$$\lim_{n \rightarrow \infty} \int_{\mathbb{R}^N} f(u_n)v_n dx = 0$$

and thus (3.15) holds.

**Claim 2.**  $\mu > 0$ .

Since  $u_n - u \rightarrow 0$  in  $L^{2+4/N}(\mathbb{R}^N)$ , we have  $\int_{\mathbb{R}^N} F(u_n - u) dx \rightarrow 0$  via Lemma 3.1 (i) and then  $\int_{\mathbb{R}^N} F(u_n) dx \rightarrow \int_{\mathbb{R}^N} F(u) dx$  by Lemma 3.2. In view of (3.11), we deduce that

$$\begin{aligned} I(u) &= \frac{1}{2} \int_{\mathbb{R}^N} |\nabla u|^2 dx - \int_{\mathbb{R}^N} F(u) dx \\ &\leq \frac{1}{2} \lim_{n \rightarrow \infty} \int_{\mathbb{R}^N} |\nabla u_n|^2 dx - \lim_{n \rightarrow \infty} \int_{\mathbb{R}^N} F(u_n) dx \\ &= \lim_{n \rightarrow \infty} I(u_n) = c < 0. \end{aligned}$$

Since, by Palais principle of symmetric criticality [19] and Pohožaev identity,

$$P(u) := \frac{N-2}{2N} \int_{\mathbb{R}^N} |\nabla u|^2 dx + \frac{1}{2} \mu \int_{\mathbb{R}^N} u^2 dx - \int_{\mathbb{R}^N} F(u) dx = 0,$$

we conclude that

$$0 > I(u) = I(u) - P(u) = \frac{1}{N} \int_{\mathbb{R}^N} |\nabla u|^2 dx - \frac{1}{2} \mu \int_{\mathbb{R}^N} u^2 dx.$$

Clearly, this implies that  $\mu > 0$ .

With Claims 1 and 2 in hand, we can now show the strong convergence. By (3.13)-(3.15) and the fact that  $\mu_n \rightarrow \mu > 0$ , we have

$$\begin{aligned} \int_{\mathbb{R}^N} |\nabla u|^2 dx + \mu \int_{\mathbb{R}^N} u^2 dx &= \int_{\mathbb{R}^N} f(u)u dx \\ &= \lim_{n \rightarrow \infty} \int_{\mathbb{R}^N} f(u_n)u_n dx = \lim_{n \rightarrow \infty} \int_{\mathbb{R}^N} |\nabla u_n|^2 dx + \mu m \\ &\geq \int_{\mathbb{R}^N} |\nabla u|^2 dx + \mu \int_{\mathbb{R}^N} u^2 dx. \end{aligned}$$

Clearly,  $\lim_{n \rightarrow \infty} \int_{\mathbb{R}^N} |\nabla u_n|^2 dx = \int_{\mathbb{R}^N} |\nabla u|^2 dx$ ,  $\int_{\mathbb{R}^N} u^2 dx = m$ , and thus  $u_n \rightarrow u$  in  $X_2$ .  $\square$

## 4 Proofs of the main results

### 4.1 Proof of Theorem 1.2

In this subsection, we shall use Theorem 2.1 to prove Theorem 1.2. Recall that  $N \geq 4$ ,  $N - 2M \neq 1$  and  $X_2 := H_{\mathcal{O}_2}^1(\mathbb{R}^N) \cap X_\tau$ . For any  $m > 0$  and  $k \in \mathbb{N}$ , we define

$$A_{m,k} := \gamma_{m,k}[\mathbb{S}^{k-1}],$$

where  $\gamma_{m,k}$  is the odd continuous mapping given by Lemma 3.4. Clearly,  $A_{m,k}$  is a closed symmetric set and  $\mathcal{G}(A_{m,k}) \geq k$  by Proposition 2.2 (iii). Therefore, the class

$$\Gamma_{m,k} := \{A \in \Sigma(S_m \cap X_2) \mid \mathcal{G}(A) \geq k\}$$

is nonempty and the minimax value

$$E_{m,k} := \inf_{A \in \Gamma_{m,k}} \sup_{u \in A} I(u)$$

is well defined.

**Lemma 4.1** (i)  $-\infty < E_{m,k} \leq E_{m,k+1} \leq 0$  for all  $m > 0$  and  $k \in \mathbb{N}$ .

(ii) For any  $k \in \mathbb{N}$ , there exists  $m(k) > 0$  large enough such that  $E_{m,k} < 0$  for all  $m > m(k)$ .

(iii) If in addition (1.2) holds, then  $E_{m,k} < 0$  for all  $m > 0$  and  $k \in \mathbb{N}$ .

(iv) For any  $k \in \mathbb{N}$ , the mapping  $m \mapsto E_{m,k}$  is nonincreasing and continuous.

**Proof.** (i) Since  $\Gamma_{m,k+1} \subset \Gamma_{m,k}$  and  $I$  is bounded from below on  $S_m \cap X_2$  by Lemma 3.1 (ii), we have

$$E_{m,k+1} \geq E_{m,k} > -\infty.$$

Let  $\gamma_{m,k}^s$  be the odd continuous mapping given by Lemma 3.4 (ii). Clearly,  $\gamma_{m,k}^s[\mathbb{S}^{k-1}] \in \Gamma_{m,k}$  and thus

$$E_{m,k} \leq \sup_{\sigma \in \mathbb{S}^{k-1}} I(\gamma_{m,k}^s[\sigma]) \quad \text{for any } s > 0. \quad (4.1)$$

In view of (3.8), we conclude that  $E_{m,k} \leq 0$ . The proof of Item (i) is complete.

(ii) This item follows from the fact that

$$E_{m,k} \leq \sup_{\sigma \in \mathbb{S}^{k-1}} I(\gamma_{m,k}[\sigma])$$

and Lemma 3.4 (i).

(iii) This item is a direct consequence of (4.1) and (3.9).

(iv) Fix  $k \in \mathbb{N}$ . To prove the claim that the mapping  $m \mapsto E_{m,k}$  is nonincreasing, we only need to show that, when  $s > m > 0$ ,

$$E_{s,k} \leq E_{m,k} + \varepsilon \quad \text{for any } \varepsilon > 0 \text{ sufficiently small.} \quad (4.2)$$

Clearly, (4.2) follows from Item (i) if  $E_{m,k} = 0$ . Thus, without loss of generality, we may assume that  $E_{m,k} < 0$ . Let  $\varepsilon \in (0, -E_{m,k}/2)$  be arbitrary. By the definition of  $E_{m,k}$ , there exists  $A_{m,k}(\varepsilon) \in \Gamma_{m,k}$  such that

$$\sup_{u \in A_{m,k}(\varepsilon)} I(u) \leq E_{m,k} + \varepsilon < 0. \quad (4.3)$$

Let

$$B_{s,k} := \{v(\cdot) = u(\cdot/t) \mid u \in A_{m,k}(\varepsilon)\},$$

where  $t := (s/m)^{1/N} > 1$ . It is clear that  $B_{s,k} \in \Gamma_{s,k}$  and thus

$$E_{s,k} \leq \sup_{v \in B_{s,k}} I(v) = \sup_{u \in A_{m,k}(\varepsilon)} I(u(\cdot/t)). \quad (4.4)$$

For any  $u \in A_{m,k}(\varepsilon)$ , by (4.3) and the fact that  $t > 1$ , we have

$$I(u(\cdot/t)) = t^N I(u) + \frac{1}{2} t^{N-2} (1-t^2) \int_{\mathbb{R}^N} |\nabla u|^2 dx \leq I(u) \leq E_{m,k} + \varepsilon.$$

In view of (4.4), we get the desired conclusion (4.2) and thus  $E_{m,k}$  is nonincreasing in  $m > 0$ .

We next show that the mapping  $m \mapsto E_{m,k}$  is continuous. Let  $s \in (0, m/2)$ . Since  $E_{m,k}$  is nonincreasing in  $m > 0$ , we see that  $E_{m-s,k}$  and  $E_{m+s,k}$  are monotonic and bounded as  $s \rightarrow 0^+$ . Therefore,  $E_{m-s,k}$  and  $E_{m+s,k}$  have limits as  $s \rightarrow 0^+$ . Noting that  $E_{m-s,k} \geq E_{m,k} \geq E_{m+s,k}$  for all  $s \in (0, m/2)$ , we have

$$\lim_{s \rightarrow 0^+} E_{m-s,k} \geq E_{m,k} \geq \lim_{s \rightarrow 0^+} E_{m+s,k}.$$

To complete the proof, we only need to prove the reverse inequality, that is,

$$\lim_{s \rightarrow 0^+} E_{m-s,k} \leq E_{m,k} \leq \lim_{s \rightarrow 0^+} E_{m+s,k}.$$

**Claim 1.**  $\lim_{s \rightarrow 0^+} E_{m-s,k} \leq E_{m,k}$ .

Let  $\varepsilon \in (0, 1)$  be arbitrary. By the definition of  $E_{m,k}$  and Item (i), there exists  $A_{m,k}(\varepsilon) \in \Gamma_{m,k}$  such that

$$\sup_{u \in A_{m,k}(\varepsilon)} I(u) \leq E_{m,k} + \varepsilon \leq 1.$$

Since  $A := \cup_{0 < \varepsilon < 1} A_{m,k}(\varepsilon) \subset S_m \cap X_2$ , we know from Lemma 3.1 (ii) that  $A$  is a bounded set in  $X_2$ . We define

$$B_{m-s,k} := \{v(\cdot) = u(\cdot/t_s) \mid u \in A_{m,k}(\varepsilon)\},$$

where  $t_s := [(m-s)/m]^{1/N} > 0$ . Clearly,  $B_{m-s,k} \in \Gamma_{m-s,k}$  and thus

$$\begin{aligned} E_{m-s,k} &\leq \sup_{v \in B_{m-s,k}} I(v) = \sup_{u \in A_{m,k}(\varepsilon)} I(u(\cdot/t_s)) \\ &\leq \sup_{u \in A_{m,k}(\varepsilon)} I(u) + \sup_{u \in A_{m,k}(\varepsilon)} |I(u(\cdot/t_s)) - I(u)| \\ &\leq E_{m,k} + \varepsilon + \sup_{u \in A} |I(u(\cdot/t_s)) - I(u)|. \end{aligned}$$

Since  $\varepsilon \in (0, 1)$  is arbitrary and  $\sup_{u \in A} |I(u(\cdot/t_s)) - I(u)|$  is independent of  $\varepsilon \in (0, 1)$ , one will obtain Claim 1 if

$$\lim_{s \rightarrow 0^+} \left( \sup_{u \in A} |I(u(\cdot/t_s)) - I(u)| \right) = 0. \quad (4.5)$$

We now prove (4.5). Noting that  $t_s$  is only dependent on  $s$ , we have

$$\sup_{u \in A} |I(u(\cdot/t_s)) - I(u)| \leq \frac{1}{2} |t_s^{N-2} - 1| \sup_{u \in A} \int_{\mathbb{R}^N} |\nabla u|^2 dx + |t_s^N - 1| \sup_{u \in A} \int_{\mathbb{R}^N} |F(u)| dx.$$

Since  $A$  is bounded in  $X_2$ , by (f1) – (f3), we see that  $\sup_{u \in A} \int_{\mathbb{R}^N} |F(u)| dx$  is bounded uniformly in  $s \in (0, m/2)$ . In view of the fact that  $\lim_{s \rightarrow 0^+} t_s = 1$ , we obtain (4.5).

**Claim 2.**  $\lim_{s \rightarrow 0^+} E_{m+s,k} \geq E_{m,k}$ .

For any  $s \in (0, m/2)$ , by the definition of  $E_{m+s,k}$ , there exists  $A_{m+s,k} \in \Gamma_{m+s,k}$  such that

$$\sup_{u \in A_{m+s,k}} I(u) \leq E_{m+s,k} + s.$$

Let  $A := \cup_{0 < s < m/2} A_{m+s,k}$ . Since  $\sup_{u \in A} \|u\|_{L^2(\mathbb{R}^N)}^2 \leq 3m/2$  and  $\sup_{u \in A} I(u) \leq m/2$  by Item (i), we know from Lemma 3.1 (ii) that  $A$  is a bounded set in  $X_2$ . Define

$$B_{m,k}(s) := \{v(\cdot) = u(\cdot/t_s) \mid u \in A_{m+s,k}\},$$



where  $t_s := [m/(m+s)]^{1/N} > 0$ . Clearly,  $B_{m,k}(s) \in \Gamma_{m,k}$  and thus

$$\begin{aligned} E_{m,k} &\leq \sup_{v \in B_{m,k}(s)} I(v) = \sup_{u \in A_{m+s,k}} I(u(\cdot/t_s)) \\ &\leq \sup_{u \in A_{m+s,k}} I(u) + \sup_{u \in A_{m+s,k}} |I(u(\cdot/t_s)) - I(u)| \\ &\leq E_{m+s,k} + s + \sup_{u \in A} |I(u(\cdot/t_s)) - I(u)|. \end{aligned}$$

Arguing as the proof of (4.5), we have that

$$\lim_{s \rightarrow 0^+} \left( \sup_{u \in A} |I(u(\cdot/t_s)) - I(u)| \right) = 0.$$

Therefore,

$$E_{m,k} \leq \lim_{s \rightarrow 0^+} \left( E_{m+s,k} + s + \sup_{u \in A} |I(u(\cdot/t_s)) - I(u)| \right) = \lim_{s \rightarrow 0^+} E_{m+s,k}.$$

The proof of Claim 2 is complete.  $\square$

**Proof of Theorem 1.2.** Clearly, Theorem 1.2 (i) and (ii) are Lemma 4.1 (i) and (iv) respectively. Let  $\mathcal{E} := X_2$  and  $\mathcal{H} := L^2(\mathbb{R}^N)$ . For any  $k \in \mathbb{N}$ , we define

$$m_k := \inf\{m > 0 \mid E_{m,k} < 0\}.$$

By Lemma 4.1 (i), (ii) and (iv), it follows that  $m_k \in [0, \infty)$ ,

$$E_{m,k} = 0 \quad \text{if } 0 < m \leq m_k, \quad E_{m,k} < 0 \quad \text{when } m > m_k.$$

Fixing  $k \in \mathbb{N}$ , when  $m > m_k$ , we have

$$-\infty < E_{m,1} \leq E_{m,2} \leq \cdots \leq E_{m,k} < 0.$$

In view of Lemma 3.1 (ii), Lemma 3.5, and Theorem 2.1 (i) and (ii), we know that  $I|_{S_m \cap X_2}$  has at least  $k$  distinct critical points associated to the levels  $E_{m,j}$  ( $j = 1, 2, \dots, k$ ). Thus, by Palais principle of symmetric criticality, we obtain Theorem 1.2 (iii). If (1.2) holds, by Lemma 4.1 (iii), we see that  $E_{m,k} < 0$  for any  $m > 0$  and  $k \in \mathbb{N}$  (and thus  $m_k = 0$  for any  $k \in \mathbb{N}$ ). Applying Theorem 2.1 (i) and (iii) to  $I|_{S_m \cap X_2}$ , we get Theorem 1.2 (iv).  $\square$

## 4.2 Proof of Theorem 1.1

This subsection is devoted to the proof of Theorem 1.1. Recall that  $N \geq 4$ ,  $X_1 := H_{O_1}^1(\mathbb{R}^N) \cap X_\tau$  and

$$E_{1,m} := \inf_{u \in S_m \cap X_1} I(u).$$

Clearly,  $E_{1,m} > -\infty$  by Lemma 3.1 (ii). Since  $X_2 \subset X_1$ , using Lemma 3.4 and arguing as the proof of Lemma 4.1, we also have

**Lemma 4.2** (i)  $-\infty < E_{1,m} \leq 0$  for all  $m > 0$ .

(ii) There exists  $m_0 > 0$  large enough such that  $E_{1,m} < 0$  for all  $m > m_0$ .

(iii) If in addition (1.2) holds, then  $E_{1,m} < 0$  for all  $m > 0$ .

(iv) The mapping  $m \mapsto E_{1,m}$  is nonincreasing and continuous.

The following two lemmas are motivated respectively by [21, Lemma 2.3 (ii)] and [21, Lemma 3.2] (see also [13, Lemma 3.2]).

**Lemma 4.3** *When  $N - 2M \neq 0$ , we have  $E_{1,m+s} \leq E_{1,m} + E_{1,s}$  for any  $m, s > 0$ .*

**Proof.** Fix  $\varepsilon > 0$ . By the definition of  $E_{1,m}$  and  $E_{1,s}$ , there exist  $u \in S_m \cap X_1 \cap C_0^\infty(\mathbb{R}^N)$  and  $v \in S_s \cap X_1 \cap C_0^\infty(\mathbb{R}^N)$  such that

$$I(u) \leq E_{1,m} + \varepsilon \quad \text{and} \quad I(v) \leq E_{1,s} + \varepsilon.$$

Since  $u$  and  $v$  have compact supports, up to a suitable translation in  $\{0\} \times \{0\} \times \mathbb{R}^{N-2M}$ , we may assume that  $\text{supp}(u) \cap \text{supp}(v) = \emptyset$ . Therefore,  $u + v \in S_{m+s} \cap X_1$  and then

$$E_{1,m+s} \leq I(u + v) = I(u) + I(v) \leq E_{1,m} + E_{1,s} + 2\varepsilon.$$

Since  $\varepsilon > 0$  is arbitrary, we obtain  $E_{1,m+s} \leq E_{1,m} + E_{1,s}$ .  $\square$

**Lemma 4.4** (i) *If  $E_{1,m}$  is achieved for some  $m > 0$ , then  $E_{1,s} < E_{1,m}$  for any  $s > m$ .*

(ii) *If  $E_{1,m}, E_{1,s}$  are achieved for some  $m, s > 0$  respectively, then  $E_{1,m+s} < E_{1,m} + E_{1,s}$ .*

**Proof.** (i) Assume that  $E_{1,m}$  is achieved at some  $u \in S_m \cap X_1$ . For any  $s > m$ , we set  $t := s/m > 1$  and  $w(x) := u(t^{-1/N}x)$ . Clearly,  $w \in S_s \cap X_1$  and

$$I(w) = tI(u) + \frac{1}{2}t^{\frac{N-2}{N}} \left(1 - t^{\frac{2}{N}}\right) \int_{\mathbb{R}^N} |\nabla u|^2 dx < tI(u) = tE_{1,m}.$$

Since  $E_{1,m} \leq 0$  by Lemma 4.2 (i), we have  $E_{1,s} \leq I(w) < tE_{1,m} \leq E_{1,m}$ .

(ii) By the assumption of Item (ii) and the argument above, we have

$$E_{1,tm} < tE_{1,m} \quad \text{for any } t > 1, \quad E_{1,\tau s} \leq \tau E_{1,s} \quad \text{for any } \tau \geq 1.$$

With loss of generality, we may assume that  $0 < s \leq m$ . Taking  $t := (m + s)/m$  and  $\tau := m/s$ , we obtain

$$E_{1,m+s} < \frac{m+s}{m}E_{1,m} = E_{1,m} + \frac{s}{m}E_{1,m} \leq E_{1,m} + E_{1,s}.$$

This completes the proof.  $\square$

**Proof of Theorem 1.1.** We define

$$m^* := \inf\{m > 0 \mid E_{1,m} < 0\}. \quad (4.6)$$

By Lemma 4.2, it is clear that  $m^* \in [0, \infty)$ ,

$$E_{1,m} = 0 \quad \text{if } 0 < m \leq m^*, \quad E_{1,m} < 0 \quad \text{when } m > m^*; \quad (4.7)$$

in particular,  $m^* = 0$  if (1.2) holds. Let us show that if  $0 < m < m^*$ , then  $E_{1,m} = 0$  is not achieved. Indeed, assuming by contradiction that  $E_{1,m}$  is achieved for some  $m \in (0, m^*)$ , in view of Lemma 4.4 (i), we have  $E_{1,m^*} < E_{1,m} = 0$  which leads to a contradiction since  $E_{1,m^*} = 0$ .

To complete the proof of Theorem 1.1, the only remaining task is to show that the infimum  $E_{1,m}$  is achieved when  $m > m^*$ . When  $N - 2M = 0$ , we have  $X_1 = X_2$  (with  $N - 2M \neq 1$ ). Since in that case  $E_{1,m} = E_{m,1}$  and  $m^* = m_1$ , the result follows directly from the property, established in Theorem 1.2, that  $E_{m,1}$  is a critical value. The rest of the proof is devoted to deal with the most delicate case, that is when  $N - 2M \neq 0$ . We shall borrow some arguments from [21, Proof of Theorem 1.2].

Fix  $m > m^*$  and let  $\{u_n\} \subset S_m \cap X_1$  be a minimizing sequence with respect to  $E_{1,m}$ . Clearly,  $\{u_n\}$  is bounded in  $X_1$  by Lemma 3.1 (ii). Up to a subsequence, we may assume that  $\lim_{n \rightarrow \infty} \int_{\mathbb{R}^N} |\nabla u_n|^2 dx$  and  $\lim_{n \rightarrow \infty} \int_{\mathbb{R}^N} F(u_n) dx$  exist. Since  $E_{1,m} < 0$  by (4.7), we have that

$$\lim_{r \rightarrow \infty} \left( \lim_{n \rightarrow \infty} \sup_{y \in \{0\} \times \{0\} \times \mathbb{R}^{N-2M}} \int_{B(y,r)} |u_n|^2 dx \right) > 0. \quad (4.8)$$

Indeed, if (4.8) does not hold, then  $u_n \rightarrow 0$  in  $L^{2+4/N}(\mathbb{R}^N)$  by Lemma 3.3 and thus

$$\lim_{n \rightarrow \infty} \int_{\mathbb{R}^N} F(u_n) dx = 0$$

via Lemma 3.1 (i); since  $I(u_n) \geq - \int_{\mathbb{R}^N} F(u_n) dx$ , a contradiction is obtained as follows:

$$0 > E_{1,m} = \lim_{n \rightarrow \infty} I(u_n) \geq - \lim_{n \rightarrow \infty} \int_{\mathbb{R}^N} F(u_n) dx = 0.$$

With (4.8) in hand, we see that there exist  $r_0 > 0$  and  $\{y_n\} \subset \{0\} \times \{0\} \times \mathbb{R}^{N-2M}$  such that

$$\lim_{n \rightarrow \infty} \int_{B(y_n, r_0)} |u_n|^2 dx > 0. \quad (4.9)$$

Since  $\{u_n(\cdot - y_n)\} \subset S_m \cap X_1$  is bounded, up to a subsequence, we may assume that  $u_n(\cdot - y_n) \rightharpoonup u$  in  $X_1$  for some  $u \in X_1$ ,  $u_n(\cdot - y_n) \rightarrow u$  in  $L^2_{\text{loc}}(\mathbb{R}^N)$  and  $u_n(\cdot - y_n) \rightarrow u$  almost everywhere in  $\mathbb{R}^N$ . Clearly,  $u \neq 0$  by (4.9). Let  $v_n := u_n(\cdot - y_n) - u$ . Noting that  $v_n \rightharpoonup 0$  in  $X_1$ , we have

$$\int_{\mathbb{R}^N} |u + v_n|^2 dx = \int_{\mathbb{R}^N} |u|^2 dx + \int_{\mathbb{R}^N} |v_n|^2 dx + o_n(1)$$

and

$$\int_{\mathbb{R}^N} |\nabla(u + v_n)|^2 dx = \int_{\mathbb{R}^N} |\nabla u|^2 dx + \int_{\mathbb{R}^N} |\nabla v_n|^2 dx + o_n(1),$$

where  $o_n(1) \rightarrow 0$  as  $n \rightarrow \infty$ . With the aid of Lemma 3.2, we also have

$$\lim_{n \rightarrow \infty} \int_{\mathbb{R}^N} F(u + v_n) dx = \int_{\mathbb{R}^N} F(u) dx + \lim_{n \rightarrow \infty} \int_{\mathbb{R}^N} F(v_n) dx.$$

Since  $I(u_n) = I(u_n(\cdot - y_n)) = I(u + v_n)$ , it follows that

$$m = \|u\|_{L^2(\mathbb{R}^N)}^2 + \lim_{n \rightarrow \infty} \|v_n\|_{L^2(\mathbb{R}^N)}^2 \quad (4.10)$$

and

$$E_{1,m} = \lim_{n \rightarrow \infty} I(u + v_n) = I(u) + \lim_{n \rightarrow \infty} I(v_n). \quad (4.11)$$

We prove below two claims and then conclude the proof.

**Claim 1.**

$$\lim_{r \rightarrow \infty} \left( \lim_{n \rightarrow \infty} \sup_{y \in \{0\} \times \{0\} \times \mathbb{R}^{N-2M}} \int_{B(y,r)} |v_n|^2 dx \right) = 0. \quad (4.12)$$

We assume by contradiction that (4.12) does not hold. Since  $\{v_n\}$  is bounded in  $X_1$ , similarly as above, up to a subsequence, there exist  $\{z_n\} \subset \{0\} \times \{0\} \times \mathbb{R}^{N-2M}$  and  $v \in X_1 \setminus \{0\}$  such that  $v_n(\cdot - z_n) \rightharpoonup v$  in  $X_1$ ,  $v_n(\cdot - z_n) \rightarrow v$  in  $L^2_{\text{loc}}(\mathbb{R}^N)$  and  $v_n(\cdot - z_n) \rightarrow v$  almost everywhere in  $\mathbb{R}^N$ . Let  $w_n := v_n(\cdot - z_n) - v$ . Then, arguing as above, we have

$$\lim_{n \rightarrow \infty} \|v_n\|_{L^2(\mathbb{R}^N)}^2 = \lim_{n \rightarrow \infty} \|v_n(\cdot - z_n)\|_{L^2(\mathbb{R}^N)}^2 = \|v\|_{L^2(\mathbb{R}^N)}^2 + \lim_{n \rightarrow \infty} \|w_n\|_{L^2(\mathbb{R}^N)}^2$$

and

$$\lim_{n \rightarrow \infty} I(v_n) = \lim_{n \rightarrow \infty} I(v_n(\cdot - z_n)) = I(v) + \lim_{n \rightarrow \infty} I(w_n).$$

In view of (4.10) and (4.11), we conclude that

$$m = \|u\|_{L^2(\mathbb{R}^N)}^2 + \|v\|_{L^2(\mathbb{R}^N)}^2 + \lim_{n \rightarrow \infty} \|w_n\|_{L^2(\mathbb{R}^N)}^2 \quad (4.13)$$

and

$$E_{1,m} = I(u) + I(v) + \lim_{n \rightarrow \infty} I(w_n). \quad (4.14)$$

Let  $s := \|u\|_{L^2(\mathbb{R}^N)}^2 > 0$ ,  $t := \|v\|_{L^2(\mathbb{R}^N)}^2 > 0$  and  $\delta := m - s - t$ . By (4.13), it is clear that

$$\delta = \lim_{n \rightarrow \infty} \|w_n\|_{L^2(\mathbb{R}^N)}^2 \geq 0.$$

To derive a contradiction, we distinguish two cases: *non-vanishing* and *vanishing*.

- *Non-vanishing: that is  $\delta > 0$ .* Let

$$t_n := \left( \frac{\delta}{\|w_n\|_{L^2(\mathbb{R}^N)}^2} \right)^{\frac{1}{N}} \quad \text{and} \quad \tilde{w}_n := w_n(\cdot/t_n).$$

Clearly,  $\lim_{n \rightarrow \infty} t_n = 1$  and  $\{\tilde{w}_n\} \subset S_\delta \cap X_1$ . Noting that

$$|I(\tilde{w}_n) - I(w_n)| \leq \frac{1}{2} |t_n^{N-2} - 1| \int_{\mathbb{R}^N} |\nabla w_n|^2 dx + |t_n^N - 1| \int_{\mathbb{R}^N} |F(w_n)| dx,$$

and that  $\int_{\mathbb{R}^N} |\nabla w_n|^2 dx$  and  $\int_{\mathbb{R}^N} |F(w_n)| dx$  are bounded, we have

$$\lim_{n \rightarrow \infty} I(w_n) = \lim_{n \rightarrow \infty} I(\tilde{w}_n) \geq E_{1,\delta}.$$

Since  $u \in S_s \cap X_1$  and  $v \in S_t \cap X_1$ , by (4.14) and Lemma 4.3, we deduce that

$$\begin{aligned} E_{1,m} &\geq I(u) + I(v) + E_{1,\delta} \\ &\geq E_{1,s} + E_{1,t} + E_{1,\delta} \geq E_{1,s+t} + E_{1,\delta} \geq E_{1,s+t+\delta} = E_{1,m}. \end{aligned}$$

Hence,  $E_{1,s}$  and  $E_{1,t}$  are achieved at  $u$  and  $v$  respectively, and

$$E_{1,s+t} = E_{1,s} + E_{1,t}. \quad (4.15)$$

By the former and Lemma 4.4 (i), we obtain  $E_{1,s+t} < E_{1,s} + E_{1,t}$  which contradicts (4.15).

• *Vanishing: that is  $\delta = 0$ .* In this case, we have  $m = s + t$  and  $\lim_{n \rightarrow \infty} \|w_n\|_{L^2(\mathbb{R}^N)} = 0$ . By Lemma 3.1 (i), it follows that  $\int_{\mathbb{R}^N} F(w_n) dx \rightarrow 0$  and thus

$$\lim_{n \rightarrow \infty} I(w_n) \geq 0.$$

Noting that  $u \in S_s \cap X_1$  and  $v \in S_t \cap X_1$ , by (4.14) and Lemma 4.3, we obtain

$$E_{1,m} \geq I(u) + I(v) \geq E_{1,s} + E_{1,t} \geq E_{1,s+t} = E_{1,m}.$$

Therefore,  $E_{1,s}$  and  $E_{1,t}$  are achieved at  $u$  and  $v$  respectively, and (4.15) holds. But, by the former and Lemma 4.4 (i), we have  $E_{1,s+t} < E_{1,s} + E_{1,t}$  which contradicts (4.15). The proof of Claim 1 is now complete.

**Claim 2.**  $\lim_{n \rightarrow \infty} \|v_n\|_{L^2(\mathbb{R}^N)}^2 = 0$ . In particular, by (4.10),  $\|u\|_{L^2(\mathbb{R}^N)}^2 = m$ .

First, by Claim 1 and Lemma 3.3, we have

$$\lim_{n \rightarrow \infty} \int_{\mathbb{R}^N} F(v_n) dx = 0. \quad (4.16)$$

If  $\lim_{n \rightarrow \infty} \|v_n\|_{L^2(\mathbb{R}^N)}^2 > 0$ , then (4.10) implies that  $s := \|u\|_{L^2(\mathbb{R}^N)}^2 < m$ . Since  $\lim_{n \rightarrow \infty} I(v_n) \geq 0$  by (4.16), in view of (4.11) and Lemma 4.2 (iv), we deduce that

$$E_{1,m} \geq I(u) \geq E_{1,s} \geq E_{1,m}.$$

Thus,  $E_{1,s}$  is achieved at  $u$  and  $E_{1,m} = E_{1,s}$ . But, by Lemma 4.4 (i), we have  $E_{1,m} < E_{1,s}$  which leads a contradiction.

**Conclusion.** Clearly,  $u \in S_m \cap X_1$  by Claim 2 and thus  $I(u) \geq E_{1,m}$ . In view of (4.11), we have

$$E_{1,m} \geq E_{1,m} + \lim_{n \rightarrow \infty} I(v_n).$$

That is,  $\lim_{n \rightarrow \infty} I(v_n) \leq 0$ . With the aid of (4.16), it then follows that

$$\frac{1}{2} \lim_{n \rightarrow \infty} \int_{\mathbb{R}^N} |\nabla v_n|^2 dx = \lim_{n \rightarrow \infty} I(v_n) + \lim_{n \rightarrow \infty} \int_{\mathbb{R}^N} F(v_n) dx \leq 0.$$

By Claim 2 again, we have  $\lim_{n \rightarrow \infty} \|v_n\|_{X_1} = 0$ . Therefore,  $u_n(\cdot - y_n) \rightarrow u$  in  $X_1$  and  $E_{1,m}$  is achieved at  $u \in S_m \cap X_1$ . We complete the proof of Theorem 1.1.  $\square$

**Remark 4.1** *In the proof of Theorem 1.1 we define the number  $m^*$  via (4.6). When we do not have (1.2), this number can be positive. To see this we assume, in addition to (f1) – (f5), that*

$$\limsup_{t \rightarrow 0} \frac{F(t)}{|t|^{2+\frac{4}{N}}} < +\infty. \quad (4.17)$$

*Since there exists  $C(f) > 0$  such that  $F(t) \leq C(f)|t|^{2+4/N}$  for any  $t \in \mathbb{R}$ , by Gagliardo-Nirenberg inequality, it follows that*

$$\int_{\mathbb{R}^N} F(u) dx \leq C(f) \int_{\mathbb{R}^N} |u|^{2+4/N} dx \leq C(f)C(N)m^{2/N} \int_{\mathbb{R}^N} |\nabla u|^2 dx$$

*for all  $u \in S_m$ . Then, for any  $m > 0$  small enough such that  $C(f)C(N)m^{2/N} \leq 1/4$ , we have*

$$I(u) \geq \frac{1}{2} \int_{\mathbb{R}^N} |\nabla u|^2 dx - \frac{1}{4} \int_{\mathbb{R}^N} |\nabla u|^2 dx = \frac{1}{4} \int_{\mathbb{R}^N} |\nabla u|^2 dx > 0.$$

*Clearly, this implies that  $E_{1,m} \geq 0$  when  $m > 0$  is small. We remark that the condition (4.17) was first introduced in [21] to show that  $m_* > 0$  may occur.*

## 5 Another proof for Theorem B

Using Theorem 2.1, the minimax theorem in Section 2, we can give another proof for Theorem B. Since the arguments are similar to that for Theorem 1.2, we just outline the proof. Recall that  $N \geq 2$  and  $H_r^1(\mathbb{R}^N)$  stands for the space of radially symmetric functions in  $H^1(\mathbb{R}^N)$ . Since the embedding  $H_r^1(\mathbb{R}^N) \hookrightarrow L^p(\mathbb{R}^N)$  is compact for any  $2 < p \leq 2+4/N$ , modifying the proof of Lemma 3.5 accordingly, we have the following compactness result.

**Lemma 5.1** *The constrained functional  $I_{|S_m \cap H_r^1(\mathbb{R}^N)}$  satisfies the condition  $(PS)_c$  for all  $c < 0$ .*

Fix  $m > 0$  and  $k \in \mathbb{N}$ . By [6, Theorem 10], there exists an odd continuous mapping  $\bar{\pi}_k : \mathbb{S}^{k-1} \rightarrow H_r^1(\mathbb{R}^N) \setminus \{0\}$  such that

$$\inf_{\sigma \in \mathbb{S}^{k-1}} \int_{\mathbb{R}^N} F(\bar{\pi}_k[\sigma]) dx \geq 1 \quad \text{and} \quad \sup_{\sigma \in \mathbb{S}^{k-1}} \|\bar{\pi}_k[\sigma]\|_{L^\infty(\mathbb{R}^N)} \leq \zeta.$$

Thus, an odd continuous mapping  $\bar{\gamma}_{m,k} : \mathbb{S}^{k-1} \rightarrow S_m \cap H_r^1(\mathbb{R}^N)$  can be introduced as follows:

$$\bar{\gamma}_{m,k}[\sigma](x) := \bar{\pi}_k[\sigma] \left( m^{-1/N} \cdot \|\bar{\pi}_k[\sigma]\|_{L^2(\mathbb{R}^N)}^{2/N} \cdot x \right), \quad x \in \mathbb{R}^N \text{ and } \sigma \in \mathbb{S}^{k-1}.$$

For any  $s > 0$ , we then define  $\bar{\gamma}_{m,k}^s[\sigma](x) := s^{N/2} \bar{\gamma}_{m,k}[\sigma](sx)$ . Arguing as the proof of Lemma 3.4, we see that  $\bar{\gamma}_{m,k}$  and  $\bar{\gamma}_{m,k}^s$  satisfy the following lemma.

**Lemma 5.2** (i) For any  $k \in \mathbb{N}$ , there exists  $\bar{m}(k) > 0$  large enough such that

$$\sup_{\sigma \in \mathbb{S}^{k-1}} I(\bar{\gamma}_{m,k}[\sigma]) < 0 \quad \text{for all } m > \bar{m}(k).$$

(ii) For any  $m > 0$  and  $k \in \mathbb{N}$ , we have

$$\limsup_{s \rightarrow 0^+} \left( \sup_{\sigma \in \mathbb{S}^{k-1}} I(\bar{\gamma}_{m,k}^s[\sigma]) \right) \leq 0.$$

If in addition (1.2) holds, then there exists  $s^* > 0$  small enough such that

$$\sup_{\sigma \in \mathbb{S}^{k-1}} I(\bar{\gamma}_{m,k}^s[\sigma]) < 0 \quad \text{for any } s \in (0, s^*).$$

Since  $\mathcal{G}(\bar{\gamma}_{m,k}[\mathbb{S}^{k-1}]) \geq k$ , the class

$$\bar{\Gamma}_{m,k} := \{A \in \Sigma(S_m \cap H_r^1(\mathbb{R}^N)) \mid \mathcal{G}(A) \geq k\}$$

is nonempty and the minimax value

$$\bar{E}_{m,k} := \inf_{A \in \bar{\Gamma}_{m,k}} \sup_{u \in A} I(u)$$

is well defined. With the aid of Lemma 5.2, repeating the argument of Lemma 4.1, we obtain

**Lemma 5.3** (i)  $-\infty < \bar{E}_{m,k} \leq \bar{E}_{m,k+1} \leq 0$  for all  $m > 0$  and  $k \in \mathbb{N}$ .

(ii) For any  $k \in \mathbb{N}$ , there exists  $\bar{m}(k) > 0$  large enough such that  $\bar{E}_{m,k} < 0$  for all  $m > \bar{m}(k)$ .

(iii) If in addition (1.2) holds, then  $\bar{E}_{m,k} < 0$  for all  $m > 0$  and  $k \in \mathbb{N}$ .

(iv) For any  $k \in \mathbb{N}$ , the mapping  $m \mapsto \bar{E}_{m,k}$  is nonincreasing and continuous.

**Conclusion.** Let  $\mathcal{E} := H_r^1(\mathbb{R}^N)$  and  $\mathcal{H} := L^2(\mathbb{R}^N)$ . For any  $k \in \mathbb{N}$ , define

$$\bar{m}_k := \inf\{m > 0 \mid \bar{E}_{m,k} < 0\}.$$

In view of Lemma 3.1 (ii), Lemma 5.1, Lemma 5.3 and Theorem 2.1, we obtain Theorem B.  $\square$

## References

- [1] T. Bartsch, S. De Valeriola, *Normalized solutions of nonlinear Schrödinger equations*, Arch. Math. **100** (2013) 75–83.
- [2] T. Bartsch, N. Soave, *A natural constraint approach to normalized solutions on nonlinear Schrödinger equations and systems*, J. Funct. Anal. **272** (2017) 4998–5037.
- [3] T. Bartsch, M. Willem, *Infinitely many nonradial solutions of a Euclidean scalar field equation*, J. Funct. Anal. **117** (1993) 447–460.
- [4] T. Bartsch, M. Willem, *Infinitely many radial solutions of a semilinear elliptic problem on  $\mathbb{R}^N$* , Arch. Rat. Mech. Anal. **124** (1993) 261–276.
- [5] H. Berestycki, P.L. Lions, *Nonlinear scalar field equations I: Existence of a ground state*, Arch. Rat. Mech. Anal. **82** (1983) 313–346.
- [6] H. Berestycki, P.L. Lions, *Nonlinear scalar field equations II: Existence of infinitely many solutions*, Arch. Rat. Mech. Anal. **82** (1983) 347–375.
- [7] H. Brezis, E. Lieb, *A relation between pointwise convergence of functions and convergence of functionals*, Proc. Amer. Math. Soc. **88** (1983), 486–490.
- [8] T. Cazenave, P.L. Lions, *Orbital stability of standing waves for some nonlinear Schrödinger equations*, Commun. Math. Phys. **85** (1982) 549–561.
- [9] J. Hirata, K. Tanaka, *Nonlinear scalar field equations with  $L^2$  constraint: mountain pass and symmetric mountain pass approaches*, reprint (2018), arXiv:1803.05139v1.
- [10] L. Jeanjean, *Approche minimax des solutions d’une équation semi-linéaire elliptique en l’absence de compacité*, PhD Thesis, EPFL, **1041** (1992).
- [11] L. Jeanjean, *Existence of solutions with prescribed norm for semilinear elliptic equations*, Nonlinear Anal. **28** (1997), 1633–1659.
- [12] L. Jeanjean, S.-S. Lu, *Nonlinear scalar field equations with general nonlinearity*, reprint (2018), arXiv:1807.07350v2.
- [13] L. Jeanjean, T.J. Luo, *Sharp nonexistence results of prescribed  $L^2$ -norm solutions for some class of Schrödinger-Poisson and quasi-linear equations*, Z. Angew. Math. Phys. **64** (2013), 937–954.
- [14] P.-L. Lions, *Symétrie et compacité dans les espaces de Sobolev*, J. Funct. Anal. **49** (1982) 315–344.
- [15] Z.L. Liu, Z.-Q. Wang, *Sign-changing solutions of nonlinear elliptic equations*, Front. Math. China **3** (2008) 221–238.
- [16] S. Lorca, P. Ubilla, *Symmetric and nonsymmetric solutions for an elliptic equation on  $\mathbb{R}^N$* , Nonlinear Anal. **58** (2004) 961–968.
- [17] J. Mederski, *Nonradial solutions of nonlinear scalar field equations*, reprint (2017), arXiv:1711.05711v3.
- [18] M. Musso, F. Pacard, J.C. Wei, *Finite-energy sign-changing solutions with dihedral symmetry for the stationary nonlinear Schrödinger equation*, J. Eur. Math. Soc. **14** (2012) 1923–1953.
- [19] R.S. Palais, *The principle of symmetric criticality*, Commun. Math. Phys. **69** (1979) 19–30.
- [20] P.H. Rabinowitz, *Minimax methods in critical point theory with applications to differential equations*, CBMS Regional Conference Series in Math., **65** American Mathematical Society, Providence, RI, 1986.
- [21] M. Shibata, *Stable standing waves of nonlinear Schrödinger equations with a general nonlinear term*, Manuscripta Math. **143** (2014) 221–237.
- [22] C.A. Stuart, *Bifurcation for Dirichlet problems without eigenvalues*, Proc. Lond. Math. Soc. **45** (1982) 169–192.
- [23] C.A. Stuart, *Bifurcation from the essential spectrum for some noncompact nonlinearities*, Math. Methods Appl. Sci. **11** (1989) 525–542.



- [24] A. Szulkin, *Ljusternik-Schnirelmann theory on  $C^1$ -manifold*, Ann. Inst. H. Poincaré Anal. Non Linéaire **5** (1988) 119–139.
- [25] M. Willem, *Minimax Theorems*, Birkhäuser Verlag (1996).