

THE NON-LINEAR GEOMETRY OF BANACH SPACES AFTER NIGEL KALTON

G. GODEFROY, G. LANCIEN AND V. ZIZLER

Dedicated to the memory of Nigel J. Kalton

ABSTRACT. This is a survey of some of the results which were obtained in the last 12 years on the non-linear geometry of Banach spaces. We focus on the contribution of the late Nigel Kalton.

1. Introduction. Four articles among Nigel Kalton's last papers are devoted to the non-linear geometry of Banach spaces ([**58**, **59**, **60**, **61**]). Needless to say, each of these works is important for the results and also for the open problems it contains. These articles followed a number of contributions due to Nigel Kalton (sometimes assisted by co-authors) which reshaped the non-linear geometry of Banach spaces during the last decade. Most of these contributions took place after Benyamini-Lindenstrauss's authoritative book [**13**] was released, and it seems that they are not yet accessible in a unified and organized way. The present survey addresses this need, in order to facilitate the access to Kalton's results (and related ones) and to help trigger further research in this widely open field of research. Nigel Kalton cannot be replaced, neither as a friend nor as the giant of mathematics he was. But his wish certainly was that research should go on, no matter what. This work is a modest attempt to fulfill this wish, and to honor his memory.

Let us outline the contents of this article. Section 2 gathers several tables, whose purpose is to present in a handy way what is known so far about the stability of several isomorphic classes under non-linear isomorphisms or embeddings. We hope that this section will provide the reader with easy access to the state of the art. Of

2010 AMS *Mathematics subject classification*. Primary 46B80, Secondary 46B20, 46B85.

The second author was partially supported by the P.H.C. Barrande 26516YG.

Received by the editors on May 15, 2012.

course, these tables contain a number of question marks, since our present knowledge is far from complete, even for classical Banach spaces. Section 3 displays several results illustrating the non-trivial fact that asymptotic structures are somewhat invariant under non-linear isomorphisms. Section 4 deals with embeddings of special graphs into Banach spaces and the use of such embeddings for showing the stability of certain properties under non-linear isomorphisms. The non-separable theory is addressed in Section 5. Non-separable spaces behave quite differently from separable ones and this promptly yields to open problems (and even to undecidable ones in ZFC). Section 6 displays the link between coarse embeddings of discrete groups (more generally of locally finite metric spaces) into the Hilbert space or super-reflexive spaces, and the classification of manifolds up to homotopy equivalence. This section attempts to provide the reader with some feeling on what the Novikov conjecture is about, and some connections between the non-linear geometry of Banach spaces and the “geometry of groups” in the sense of Gromov. Finally, Section 7 is devoted to the Lipschitz-free spaces associated with a metric space, their use and their structure (or at least, what is known about it). Sections 2 and 6 contain no proof, but other sections do. These proofs were chosen in order to provide information on the tools we need.

This work is a survey, but it contains some statements (such as Theorems 3.8 and 3.12 or the last Remark in Section 5) which were not published before. Each section contains a number of commented open questions. It is interesting to observe that many of these questions are fairly simple to state. Answering them could be less simple. Our survey demonstrates that non-linear geometry of Banach spaces is a meeting point for a variety of techniques, which have to join forces in order to allow progress. It is our hope that the present work will help stimulate such efforts. We should, however, make it clear that our outline of Nigel Kalton’s last papers does not exhaust the content of these articles. We strongly advise the interested reader to consult them for her/his own research.

2. Tables. This section consists of five tables: Table 1 lists a number of classical spaces and checks when these Banach spaces are characterized by their metric or their uniform structure. Table 2 displays what is known about Lipschitz embeddings from a classical Banach space into

another, and Table 3 does the same for uniform embeddings. Table 4 investigates the stability of certain isomorphism classes (relevant to a classical property) under Lipschitz or uniform homeomorphism. And finally, Table 5 does the same for non-linear embeddability.

References are given within the tables themselves, but in order to improve readability, we almost always use symbols (whose meaning is explained below) rather than using the numbering of the reference list. Question marks mean, of course, that to the best of our knowledge, the corresponding question is still open.

Our notation for Banach spaces is standard. All Banach spaces will be real. From the recent textbooks that may be used in the area we mention [5, 32].

♣ = Benyamini-Lindenstrauss book [13].

♠ = Kalton recent papers.

‡ = Mendel-Naor papers [76] and [78].

△ = Godefroy-Kalton-Lancien papers [38] and [39].

□ = Godefroy-Kalton paper on free spaces [37].

♡ = Johnson-Lindenstrauss-Schechtman paper [51].

◇ = Textbook [32].

◊ = Basic linear theory or topology.

? = Unknown to the authors.

We say that a Banach space X is determined by its Lipschitz (respectively uniform) structure if a Banach space Y is linearly isomorphic to X whenever Y is Lipschitz homeomorphic (respectively uniformly homeomorphic) to X .

A Lipschitz embedding of a Banach space X into a Banach space Y is a Lipschitz homeomorphism from X onto a *subset* (in general non linear) of Y .

A uniform embedding of a Banach space X into a Banach space Y is a uniform homeomorphism from X onto a *subset* of Y . Let us also

TABLE 1. Spaces determined by weaker structures.

Space	Determined by its Lipschitz Structure	Determined by its uniform Structure	
ℓ_2	yes	yes	♣
$1 < p < \infty$ $p \neq 2$	yes	yes	♣
ℓ_1	?	?	
c_0	yes	?	
L_p $1 < p < \infty$ $p \neq 2$	yes	?	
L_1	♣		
$C[0, 1]$?	?	
$\ell_2(c)$	yes	yes	♣
$c_0(c)$	no	no	♣
ℓ_∞	?	?	
$\ell_p \oplus \ell_q$ $1 < p < q < \infty$ $p, q \neq 2$	yes	yes	♣+♣
$\ell_p \oplus \ell_2$ $1 < p < \infty$ $p \neq 2$	yes	?	
J	♣		
James' space	yes	?	
	♣+[22]		

TABLE 2. Lipschitz embeddings from the 1st column into the 1st row.

Space	ℓ_2	ℓ_q $1 < q < \infty$ $q \neq 2$	ℓ_1	c_0	L_q $1 < q < \infty$ $q \neq 2$	L_1	$C[0, 1]$	$\ell_2(c)$	$c_0(c)$	ℓ_∞
ℓ_2	yes	no	no	yes	yes	yes	yes	yes	yes	yes
ℓ_p $1 < p < \infty$ $p \neq 2$	no	yes iff $p = q$	no	yes	yes iff $q \leq p < 2$ or $p = q$	yes iff $p < 2$	yes	no	yes	yes
ℓ_1	no	no	yes	yes	no	yes	yes	no	yes	yes
c_0	no	no	no	yes	no	no	yes	no	yes	yes
L_p $1 < p < \infty$ $p \neq 2$	no	no	no	yes	yes iff $q \leq p < 2$ or $p = q$	yes iff $p < 2$	yes	no	yes	yes
L_1	no	no	no	yes	no	yes	yes	no	yes	yes
$C[0, 1]$	no	no	no	yes	no	no	yes	no	yes	yes
$\ell_2(c)$	no	no	no	no	no	no	no	yes	?	yes
$c_0(c)$	no	no	no	no	no	no	no	no	yes	yes
ℓ_∞	no	no	no	no	no	no	no	no	?	yes

mention that the same table can be written about coarse embeddings (see the definition in Section 3). The set of references to be used is the same except for two papers by Nowak, who proved in [83] that, for

TABLE 3. Uniform Embeddings from the 1st column into the 1st row.

Space	ℓ_2	ℓ_q $q \in (1, \infty)$ $q \neq 2$	ℓ_1	c_0	L_q $q \in (1, \infty)$ $q \neq 2$	L_1	$C[0, 1]$	$\ell_2(c)$	$c_0(c)$	ℓ_∞
ℓ_2	yes	yes ♣	yes ♣	yes ♣	yes ♣	yes ♣	yes ♣	yes ♣	yes ♣	yes ♣
ℓ_p $p \in (1, \infty)$ $p \neq 2$	yes iff $p < 2$ ♣	yes if $p \leq q$ or $p < 2$ [4]+♣ no if $p > 2$ and $q < p$ ♣	yes iff $p < 2$ ♣	yes ♣	yes iff $p \leq q$ or $p < 2$ ♣	yes iff $p < 2$ ♣	yes ♣	yes iff $p < 2$ ♣	yes ♣	yes ♣
ℓ_1	yes ♣	yes ♣	yes ♣	yes ♣	yes ♣	yes ♣	yes ♣	yes ♣	yes ♣	yes ♣
c_0	no ♣	no ♣ or ♣	no ♣	yes ♣	no ♣ or ♣	no ♣	yes ♣	no ♣	yes ♣	yes ♣
L_p $p \in (1, \infty)$ $p \neq 2$	yes iff $p < 2$ ♣	no if $p > 2$ and $q < p$ ♣ yes if $p < 2$ ♣ ? if $2 < p \leq q$	yes iff $p < 2$ ♣	yes ♣	yes iff $p \leq q$ or $p < 2$ ♣	yes iff $p < 2$ ♣	yes ♣	yes iff $p < 2$ ♣	yes ♣	yes ♣
L_1	yes ♣	yes ♣	yes ♣	yes ♣	yes ♣	yes ♣	yes ♣	yes ♣	yes ♣	yes ♣
$C[0, 1]$	no ♣	no ♣ or ♣	no ♣	yes ♣	no ♣ or ♣	no ♣	yes ♣	no ♣	yes ♣	yes ♣
$\ell_2(c)$	no ♣	no ♣	no ♣	no ♣	no ♣	no ♣	no ♣	yes ♣	?	yes ♣
$c_0(c)$	no ♣	no ♣	no ♣	no ♣	no ♣	no ♣	no ♣	no ♣	yes ♣	yes ♣
ℓ_∞	no ♣	no ♣	no ♣	no ♣	no ♣	no ♣	no ♣	no ♣	?	yes ♣

any $p \in [1, \infty)$, ℓ_2 coarsely embeds into ℓ_p and in [84] that, for $p < 2$, L_p coarsely embeds into ℓ_2 . A larger table could mention the work of Kraus on embeddings between Orlicz sequence spaces [65].

More precisely, the question addressed is the following. If a Banach space X Lipschitz or coarse-Lipschitz (see definition in Section 3) embeds into a Banach space Y which has one of the properties listed in the first column, does X satisfy the same property?

3. Uniform and asymptotic structures of Banach spaces.

In this section we will study the stability of the uniform asymptotic smoothness or uniform asymptotic convexity of a Banach space under non linear maps such as uniform homeomorphisms and coarse-Lipschitz embeddings.

TABLE 4. Stability under type of homeomorphism.

Property	Lipschitz	uniform	
Hilbertian	yes	yes	
superreflexivity	yes	♣	
reflexivity	yes	♣	
RNP	♣	no	
Asplund	yes	♣	
containment of ℓ_1	?	no	
containment of c_0	?	♣	
BAP	yes	?	
Commuting BAP	□	no	
Existence of Schauder basis	?	♥	
Existence of M-basis	?	♥	
Existence of unconditional basis	no	no	
renorming by Frechet smooth norm	◇	no	
renorming by LUR norm	?	♣	
renorming by UG norm	no	?	
renorming by WUR norm	◇	no	
renorming by AUS norm	?	?	
	yes	yes	△

TABLE 5. Properties shared by embedded spaces.

Property	Lipschitz	coarse-Lipschitz	
Hilbertian	yes	yes	
superreflexivity	♣	♣	
reflexivity	yes	yes	
RNP	♣	♣	
Asplund	yes	no	
renorming by Frechet smooth norm	♣	♣	
renorming by LUR norm	no	?	
renorming by UG norm	◇	no	
renorming by WUR norm	?	?	
renorming by AUS norm	no	no	
reflexive+renorming by AUS norm	yes	?	
reflexive+renorming by AUS norm +renorming by AUC norm	♣	yes	[9]

3.1. Notation–Introduction.

Definition 3.1. Let (M, d) and (N, δ) be two metric spaces and

$f : M \rightarrow N$ be a mapping.

- (a) f is a *Lipschitz isomorphism* (or *Lipschitz homeomorphism*) if f is a bijection and f and f^{-1} are Lipschitz. We denote $M \stackrel{\text{Lip}}{\sim} N$, and we say that M and N are *Lipschitz equivalent*.
- (b) f is a *uniform homeomorphism* if f is a bijection and f and f^{-1} are uniformly continuous (we denote $M \stackrel{\text{UH}}{\sim} N$).
- (c) If (M, d) is unbounded, we define, for all $s > 0$,

$$\text{Lip}_s(f) = \sup \left\{ \frac{\delta(f(x), f(y))}{d(x, y)}, d(x, y) \geq s \right\}$$

and

$$\text{Lip}_\infty(f) = \inf_{s>0} \text{Lip}_s(f).$$

f is said to be *coarse Lipschitz* if $\text{Lip}_\infty(f) < \infty$.

- (d) f is a *coarse Lipschitz embedding* if there exist $A > 0$, $B > 0$, $\theta \geq 0$, such that, for all $x, y \in M$,

$$d(x, y) \geq \theta \implies \text{Ad}(x, y) \leq \delta(f(x), f(y)) \leq Bd(x, y).$$

We denote $M \stackrel{\text{CL}}{\hookrightarrow} N$.

More generally, f is a *coarse embedding* if there exist two real-valued functions ρ_1 and ρ_2 such that $\lim_{t \rightarrow +\infty} \rho_1(t) = +\infty$ and, for all $x, y \in M$,

$$\rho_1(d(x, y)) \leq \delta(f(x), f(y)) \leq \rho_2(d(x, y)).$$

- (e) An (a, b) -net in the metric space M is a subset \mathcal{M} of M such that, for every $z \neq z'$ in \mathcal{M} , $d(z, z') \geq a$ and, for every x in M , $d(x, \mathcal{M}) < b$.

Then a subset \mathcal{M} of M is a *net* in M if it is an (a, b) -net for some $0 < a \leq b$.

- (f) Note that two nets in the same infinite-dimensional Banach space are always Lipschitz equivalent (see [13, Proposition 10.22]).

Then two infinite-dimensional Banach spaces X and Y are said to be *net equivalent*, and we denote $X \stackrel{N}{\sim} Y$, if there exist a net \mathcal{M} in M and a net \mathcal{N} in N such that \mathcal{M} and \mathcal{N} are Lipschitz equivalent.

Remark. It follows easily from the triangle inequality that a uniformly continuous map defined on a Banach space is coarse Lipschitz, and a uniform homeomorphism between Banach spaces is a bi-coarse Lipschitz bijection (see [13, Proposition 1.11] for details). Therefore, if X and Y are uniformly homeomorphic Banach spaces, then they are net equivalent. It has been proved only recently by Kalton in [59] that there exist two net equivalent Banach spaces that are not uniformly homeomorphic. However, the finite-dimensional structures of Banach spaces are preserved under net equivalence (see [13, Proposition 10.19], or Theorem 3.3 below).

The main question addressed in this section is the problem of the uniqueness of the uniform (or net) structure of a given Banach space. In other words, whether $X \overset{UH}{\sim} Y$ (or $X \overset{N}{\sim} Y$) implies that X is linearly isomorphic to Y (which we shall denote $X \simeq Y$). Even in the separable case, the general answer is negative. Indeed, Ribe [95] proved the following.

Theorem 3.2 ([95]). *Let $(p_n)_{n=1}^\infty$ in $(1, +\infty)$ be a strictly decreasing sequence such that $\lim p_n = 1$. Denote $X = (\sum_{n=1}^\infty L_{p_n})_{\ell_2}$. Then $X \overset{UH}{\sim} X \oplus L_1$.*

Therefore, reflexivity is not preserved under coarse-Lipschitz embeddings or even uniform homeomorphisms. On the other hand, Ribe [96] proved in fact that local properties of Banach spaces are preserved under coarse-Lipschitz embeddings. More precisely,

Theorem 3.3 ([96]). *Let X and Y be two Banach spaces such that $X \overset{CL}{\hookrightarrow} Y$. Then there exists a constant $K \geq 1$ such that, for any finite-dimensional subspace E of X , there is a finite-dimensional subspace F of Y which is K -isomorphic to E .*

Remark. If we combine this result with Kwapien's theorem, we immediately obtain that a Banach space which is net equivalent to ℓ_2 is linearly isomorphic to ℓ_2 .

As announced, we will concentrate on some asymptotic properties of Banach spaces. So let us give the relevant definitions.

Definition 3.4. Let $(X, \|\cdot\|)$ be a Banach space and $t > 0$. We denote by B_X the closed unit ball of X and by S_X its unit sphere. For $x \in S_X$ and Y a closed linear subspace of X , we define

$$\bar{\rho}(t, x, Y) = \sup_{y \in S_Y} \|x + ty\| - 1$$

and

$$\bar{\delta}(t, x, Y) = \inf_{y \in S_Y} \|x + ty\| - 1.$$

Then

$$\bar{\rho}_X(t) = \sup_{x \in S_X} \inf_{\dim(X/Y) < \infty} \bar{\rho}(t, x, Y)$$

and

$$\bar{\delta}_X(t) = \inf_{x \in S_X} \sup_{\dim(X/Y) < \infty} \bar{\delta}(t, x, Y).$$

The norm $\|\cdot\|$ is said to be *asymptotically uniformly smooth* (in short, AUS) if

$$\lim_{t \rightarrow 0} \frac{\bar{\rho}_X(t)}{t} = 0.$$

It is said to be *asymptotically uniformly convex* (in short AUC) if, for all $t > 0$,

$$\bar{\delta}_X(t) > 0.$$

These moduli have been first introduced by Milman in [80]. We also refer the reader to [27, 50] for reviews on these.

Examples.

- (1) If $X = (\sum_{n=1}^{\infty} F_n)_{\ell_p}$, $1 \leq p < \infty$, and the F_n 's are finite-dimensional, then $\bar{\rho}_X(t) = \bar{\delta}_X(t) = (1 + t^p)^{1/p} - 1$. Actually, if a separable reflexive Banach space has equivalent norms with moduli of asymptotic convexity and smoothness of power type p , then it is isomorphic to a subspace of an l_p -sum of finite-dimensional spaces [50].
- (2) For all $t \in (0, 1)$, $\bar{\rho}_{c_0}(t) = 0$. And again, if X is separable and $\bar{\rho}_X(t_0) = 0$ for some $t_0 > 0$, then X is isomorphic to a subspace of c_0 [38].

We conclude this introduction by mentioning the open questions that we will comment on in the course of this section.

Problem 1. Let $1 < p < \infty$ and $p \neq 2$. Does $\ell_p \oplus \ell_2$ have a unique uniform or net structure? Does L_p have a unique uniform or net structure?

Problem 2. Assume that Y is a reflexive AUS Banach space and that X is a Banach space which coarse-Lipschitz embeds into Y . Does X admit an equivalent AUS norm?

Problem 3. Assume that Y is an AUC Banach space and that X is a Banach space which coarse-Lipschitz embeds into Y . Does X admit an equivalent AUC norm?

3.2. The approximate midpoints principle. Given a metric space X , two points $x, y \in X$, and $\delta > 0$, the approximate metric midpoint set between x and y with error δ is the set:

$$\text{Mid}(x, y, \delta) = \left\{ z \in X : \max\{d(x, z), d(y, z)\} \leq (1 + \delta) \frac{d(x, y)}{2} \right\}.$$

The use of approximate metric midpoints in the study of nonlinear geometry is due to Enflo in an unpublished paper and has since been used extensively, e.g., [17, 41, 51].

The following version of the approximate midpoint lemma was formulated in [63] (see also [13, Lemma 10.11]).

Proposition 3.5. *Let X be a normed space, and suppose M is a metric space. Let $f : X \rightarrow M$ be a coarse Lipschitz map. If $\text{Lip}_\infty(f) > 0$, then for any $t, \varepsilon > 0$ and any $0 < \delta < 1$, there exist $x, y \in X$ with $\|x - y\| > t$ and*

$$f(\text{Mid}(x, y, \delta)) \subset \text{Mid}(f(x), f(y), (1 + \varepsilon)\delta).$$

In view of this proposition, it is natural to study the approximate metric midpoints in ℓ_p . This is done in the next lemma, which is rather elementary and can be found in [63].

Lemma 3.6. *Let $1 \leq p < \infty$. We denote $(e_i)_{i=1}^{\infty}$ as the canonical basis of ℓ_p and, for $N \in \mathbb{N}$, let E_N be the closed linear span of $\{e_i, i > N\}$. Now let $x, y \in \ell_p$, $\delta \in (0, 1)$, $u = x + y/2$ and $v = x - y/2$. Then*

- (i) *There exists $N \in \mathbb{N}$ such that $u + \delta^{1/p}\|v\|B_{E_N} \subset \text{Mid}(x, y, \delta)$.*
- (ii) *There is a compact subset K of ℓ_p such that $\text{Mid}(x, y, \delta) \subset K + 2\delta^{1/p}\|v\|B_{\ell_p}$.*

We can now combine Proposition 3.5 and Lemma 3.6 to obtain:

Corollary 3.7. *Let $1 \leq p < q < \infty$. Then ℓ_q does not coarse Lipschitz embed into ℓ_p .*

Remark. This statement can be found in [63] but was implicit in [51]. It already indicates that, because of the approximate midpoint principle, some uniform asymptotic convexity has to be preserved under coarse Lipschitz embeddings. This idea will be pushed much further in Section 3.6.

3.3. Gorelik principle and applications. Our goal is now to study the stability of the uniform asymptotic smoothness under non linear maps. The first tool that we shall describe is the Gorelik principle. It was initially devised by Gorelik in [41] to prove that ℓ_p is not uniformly homeomorphic to L_p , for $1 < p < \infty$. Then it was developed by Johnson, Lindenstrauss and Schechtman [51] to prove that, for $1 < p < \infty$, ℓ_p has a unique uniform structure. It is important to underline the fact that the Gorelik principle is only valid for certain bijections. In fact, the uniqueness of the uniform structure of ℓ_p can be proved without the Gorelik principle, by using results on the embeddability of special metric graphs as we shall see in subsection 3.4. Nevertheless, some other results still need the use of the Gorelik principle. This principle is usually stated for homeomorphisms with uniformly continuous inverse (see [13, Theorem 10.12]). Although it is probably known, we have not found its version for net equivalences. Note that a Gorelik principle is proved in [13, Proposition 10.20] for net equivalences between a Banach space and ℓ_p . So we will describe here how to obtain a general statement.

Theorem 3.8 (Gorelik principle). *Let X and Y be two Banach spaces. Let X_0 be a closed linear subspace of X of finite codimension. If X and Y are net equivalent, then there are continuous maps $U : X \rightarrow Y$ and $V : Y \rightarrow X$, and constants $K, C, \alpha_0 > 0$ such that for all $x \in X$,*

$$\|VUx - x\| \leq C$$

and, for all $y \in Y$,

$$\|UVy - y\| \leq C$$

and, for all $\alpha > \alpha_0$, there is a compact subset M of Y so that

$$\frac{\alpha}{16K}B_Y \subset M + CB_Y + U(\alpha B_{X_0}).$$

Proof. Suppose that \mathcal{N} is a net of the Banach space X , that \mathcal{M} is a net of the Banach space Y and that \mathcal{N} and \mathcal{M} are Lipschitz equivalent. We will assume as we may that \mathcal{N} and \mathcal{M} are $(1, \lambda)$ -nets for some $\lambda > 1$ and that $\mathcal{N} = (x_i)_{i \in I}$, $\mathcal{M} = (y_i)_{i \in I}$ with for all $i, j \in I$,

$$K^{-1}\|x_i - x_j\| \leq \|y_i - y_j\| \leq K\|x_i - x_j\|,$$

for some $K \geq 1$. Let us denote by $B_E(x, \lambda)$ the open ball of center x and radius λ in the Banach space E . Then we can find a continuous partition of unity $(f_i)_{i \in I}$ subordinate to $(B_X(x_i, \lambda))_{i \in I}$ and a continuous partition of unity $(g_i)_{i \in I}$ subordinate to $(B_Y(y_i, \lambda))_{i \in I}$. Now we set:

$$Ux = \sum_{i \in I} f_i(x)y_i, \quad x \in X$$

and

$$Vy = \sum_{i \in I} g_i(y)x_i, \quad y \in Y.$$

The maps U and V are clearly continuous. We shall now state and prove two lemmas about them.

Lemma 3.9.

- (i) *Let $x \in X$ be such that $\|x - x_i\| \leq r$, then $\|Ux - y_i\| \leq K(\lambda + r)$.*
- (ii) *Let $y \in Y$ be such that $\|y - y_i\| \leq r$, then $\|Vy - x_i\| \leq K(\lambda + r)$.*

Proof. We will only prove (i). If $f_j(x) \neq 0$, then $\|x - x_j\| \leq \lambda$. So $\|x_i - x_j\| \leq \lambda + r$ and $\|y_i - y_j\| \leq K(\lambda + r)$. We finish the proof by writing

$$Ux - y_i = \sum_{j, f_j(x) \neq 0} f_j(x)(y_j - y_i). \quad \square$$

Lemma 3.10. *Let $C = (1 + K + 2K^2)\lambda$. Then for all $x \in X$,*

$$\|VUx - x\| \leq C$$

and for all $y \in Y$,

$$\|UVy - y\| \leq C.$$

Proof. We only need to prove one inequality. So let $x \in X$, and pick $i \in I$ such that $\|x - x_i\| \leq \lambda$. By the previous lemma, we have $\|Ux - y_i\| \leq 2K\lambda$ and $\|VUx - x_i\| \leq \lambda(K + 2K^2)$. Thus, $\|VUx - x\| \leq (1 + K + 2K^2)\lambda$. \square

We now recall the crucial ingredient in the proof of the Gorelik principle (see [13, Theorem 10.12, proof step (i)]). This statement relies on Brouwer's fixed point theorem and on the existence of Bartle-Graves continuous selectors. We refer the reader to [13] for its proof.

Proposition 3.11. *Let X_0 be a finite-codimensional subspace of X . Then, for any $\alpha > 0$, there is a compact subset A of $(\alpha/2)B_X$ such that for every continuous map $\phi : A \rightarrow X$ satisfying $\|\phi(a) - a\| \leq \alpha/4$ for all $a \in A$, we have that $\phi(A) \cap X_0 \neq \emptyset$.*

We are now ready to finish the proof of Theorem 3.8. Fix $\alpha > 0$ such that $\alpha > \max\{8C, 96K\lambda\}$ and $y \in (\alpha/16K)B_Y$, and define $\phi : A \rightarrow X$ by $\phi(a) = V(y + Ua)$. The map ϕ is clearly continuous, and we have that for all $a \in A$:

$$\begin{aligned} \|\phi(a) - a\| &\leq \|V(y + Ua) - VUa\| + \|VUa - a\| \\ &\leq \frac{\alpha}{8} + \|V(y + Ua) - VUa\|. \end{aligned}$$

Now, pick i so that $\|Ua - y_i\| \leq \lambda$ and j so that $\|y + Ua - y_j\| \leq \lambda$. Then $\|VUa - x_i\| \leq 2K\lambda$ and $\|V(y + ua) - x_j\| \leq 2K\lambda$. But,

$$\begin{aligned} \|x_i - x_j\| &\leq K\|y_i - y_j\| \\ &\leq K\|y_i - Ua\| + K\|Ua + y - y_j\| + K\|y\| \\ &\leq (2\lambda + \|y\|)K. \end{aligned}$$

So

$$\|V(y + Ua) - VUa\| \leq 6K\lambda + K\|y\| \leq 6K\lambda + \frac{\alpha}{16} \leq \frac{\alpha}{8}.$$

Thus $\|\phi(a) - a\| \leq \alpha/4$.

So it follows from Proposition 3.11 that there exists $a \in A$ such that $\phi(a) \in X_0$. Besides, $\|a\| \leq \alpha/2$ and $\|\phi(a) - a\| \leq \alpha/4$, so $\phi(a) = V(y+Ua) \in \alpha B_{X_0}$. But we have that $\|UV(y+Ua) - (y+Ua)\| \leq C$. So, if we consider the compact set $M = -U(A)$, we have that $y \in M + CB_Y + U(\alpha B_{X_0})$. This finishes the proof of Theorem 3.8. \square

We can now apply the above Gorelik principle to obtain the net version of a result that appeared in [39]. This result is new.

Theorem 3.12. *Let X and Y be Banach spaces. Assume that X is net equivalent to Y and that X is AUS. Then Y admits an equivalent AUS norm. More precisely, if $\bar{\rho}_X(t) \leq Ct^p$ for $C > 0$ and $p \in (1, \infty)$, then, for any $\varepsilon > 0$, Y admits an equivalent norm $\|\cdot\|_\varepsilon$ so that $\bar{\rho}_{\|\cdot\|_\varepsilon}(t) \leq C_\varepsilon t^{p-\varepsilon}$ for some $C_\varepsilon > 0$.*

The proof is actually done by constructing a sequence of dual norms as follows: for all $y^* \in Y^*$,

$$|y^*|_k = \sup \left\{ \frac{\langle y^*, Ux - Ux' \rangle}{\|x - x'\|}, \|x - x'\| \geq 4^k \right\}.$$

For k large enough they are all equivalent. Then, for N large enough, the predual norm of the norm defined by, for all $y^* \in Y^*$,

$$\|y^*\|_N = \frac{1}{N} \sum_{k=k_0+1}^{k=k_0+N} |y^*|_k,$$

is the dual of an equivalent AUS norm on Y with the desired modulus of asymptotic smoothness. The proof follows the lines of the argument given in [39] but uses the above version of Gorelik principle.

Remarks.

- (1) It must be pointed out that the quantitative estimate in the above result is optimal as it follows from a remarkable example obtained by Kalton in [61].
- (2) In [14] Borel-Mathurin provides explicit estimates of the Szlenk indices of Orlicz sequence spaces. This allows her to derive from Theorem 3.12 some applications to uniform homeomorphisms between subspaces and quotients of Orlicz spaces.
- (3) If the Banach spaces X and Y are Lipschitz equivalent and X is AUS, then the norm defined on Y^* by:

$$|y^*| = \sup \left\{ \frac{\langle y^*, Ux - Ux' \rangle}{\|x - x'\|}, x \neq x' \right\}$$

is a dual norm of a norm $\|\cdot\|$ on Y such that $\bar{\rho}_{\|\cdot\|}(t) \leq c\bar{\rho}_X(ct)$ for some $c > 0$. When X is a subspace of c_0 , this implies that Y is isomorphic to a subspace of c_0 . Finally, when $X = c_0$, one gets that X is isomorphic to c_0 (see [38]).

- (4) Other results were originally derived from the Gorelik principle. We have chosen to present them in the next subsections as consequences of more recent and possibly more intuitive graph techniques introduced by Kalton and Randrianarivony in [63] and later developed by Kalton in [60].

3.4. Uniform asymptotic smoothness and Kalton-Randrianarivony's graphs. The fundamental result of this section is about the minimal distortion of some special metric graphs into a reflexive and asymptotically uniformly smooth Banach space. These graphs have been introduced by Kalton and Randrianarivony in [63] and are defined as follows. Let \mathbb{M} be an infinite subset of \mathbb{N} and $k \in \mathbb{N}$, and fix $a = (a_1, \dots, a_k)$ a sequence of non zero real numbers. We denote

$$G_k(\mathbb{M}) = \{\bar{n} = (n_1, \dots, n_k), n_i \in \mathbb{M} \quad n_1 < \dots < n_k\}.$$

Then we equip $G_k(\mathbb{M})$ with the distance, for all $\bar{n}, \bar{m} \in G_k(\mathbb{M})$,

$$d_a(\bar{n}, \bar{m}) = \sum_{j, n_j \neq m_j} |a_j|.$$

Note also that it is easily checked that $\bar{\rho}_Y$ is an Orlicz function. Then, we define the Orlicz sequence space:

$$\ell_{r\text{ho}_Y} = \left\{ a \in \mathbb{R}^{\mathbb{N}}, \quad \text{there exists } r > 0 \sum_{n=1}^{\infty} \bar{\rho}_Y \left(\frac{|a_n|}{r} \right) < \infty \right\},$$

equipped with the Luxemburg norm

$$\|a\|_{\bar{\rho}_Y} = \inf \left\{ r > 0, \sum_{n=1}^{\infty} \bar{\rho}_Y \left(\frac{|a_n|}{r} \right) \leq 1 \right\}.$$

Theorem 3.13. [63]. *Let Y be a reflexive Banach space, \mathbb{M} an infinite subset of \mathbb{N} and $f : (G_k(\mathbb{M}), d_a) \rightarrow Y$ a Lipschitz map. Then, for any $\varepsilon > 0$, there exists an infinite subset \mathbb{M}' of \mathbb{M} such that:*

$$\text{diam } f(G_k(\mathbb{M}')) \leq 2e\text{Lip}(f)\|a\|_{\bar{\rho}_Y} + \varepsilon.$$

The proof is done by induction on k and uses iterated weak limits of subsequences and a Ramsey argument. Such techniques will be displayed in the next two sections.

Remark. The reflexivity assumption is important. Indeed, by Aharoni's theorem, the spaces $(G_k(\mathbb{N}), d_a)$ Lipschitz embed into c_0 with a distortion controlled by a uniform constant. But $\|a\|_{\bar{\rho}_{c_0}} = \|a\|_{\infty}$, while $\text{diam } G_k(\mathbb{M}') = \|a\|_1$.

As is described in [60], one can deduce the following.

Corollary 3.14. *Let X be a Banach space and Y a reflexive Banach space. Assume that X coarse Lipschitz embeds into Y . Then there exists $C > 0$ such that, for any normalized weakly null sequence $(x_n)_{n=1}^{\infty}$ in X and any sequence $a = (a_1, \dots, a_k)$ of non zero real numbers, there is an infinite subset \mathbb{M} of \mathbb{N} such that:*

$$\left\| \sum_{i=1}^k a_i x_{n_i} \right\| \leq C \|a\|_{\bar{\rho}_Y}, \quad \text{for every } \bar{n} \in G_k(\mathbb{M}).$$

Proof. The result is obtained by applying Theorem 3.13 to $f = g \circ h$, where g is a coarse-Lipschitz embedding from X into Y and $h : (G_k(\mathbb{N}), d_a) \rightarrow X$ is defined by $h(\bar{n}) = \lambda \sum_{i=1}^k a_i x_{n_i}$ for some large enough $\lambda > 0$. \square

In fact, this is stated in [60] in the following more abstract way.

Corollary 3.15. *Let X be a Banach space and Y a reflexive Banach space. Assume that X coarse Lipschitz embeds into Y . Then there exists $C > 0$ such that, for any spreading model $(e_i)_i$ of a normalized weakly null sequence in X (whose norm is denoted $\|\cdot\|_S$) and any finitely supported sequence $a = (a_i)$ in \mathbb{R} :*

$$\left\| \sum a_i e_i \right\|_S \leq C \|a\|_{\bar{p}_Y}.$$

3.5. Applications. The first consequence is the following.

Corollary 3.16. *Let $1 \leq q \neq p < \infty$. Then ℓ_q does not coarse Lipschitz embed into ℓ_p .*

Proof. If $q < p$, this follows immediately from the previous results. If $q > p$, this is Corollary 3.7. \square

Then we can deduce the following result, proven in [51] under the assumption of uniform equivalence.

Theorem 3.17 ([51]). *Let $1 < p < \infty$ and X be a Banach space such that $X \overset{N}{\sim} \ell_p$. Then $X \simeq \ell_p$.*

Proof. Suppose that $X \overset{N}{\sim} \ell_p$, with $1 < p < \infty$. We may assume that $p \neq 2$. Then the ultra-products $X_{\mathcal{U}}$ and $(\ell_p)_{\mathcal{U}}$ are Lipschitz isomorphic, and it follows from the classical Lipschitz theory that X is isomorphic to a complemented subspace of $L_p = L_p([0, 1])$. Now, it follows from Corollary 3.16 that X does not contain any isomorphic copy of ℓ_2 . Then we can conclude with a classical result of Johnson and Odell [52] which asserts that any infinite-dimensional complemented subspace of L_p that does not contain any isomorphic copy of ℓ_2 is isomorphic to ℓ_p . \square

Remark. These linear arguments are taken from [51]. Note that the key step was to show that X does not contain any isomorphic copy of ℓ_2 . In the original paper [51] this relied on the Gorelik principle. We have chosen to present here a proof using this graph argument. In fact, more can be deduced from this technique.

Corollary 3.18. *Let $1 \leq p < q < \infty$ and $r \geq 1$ be such that $r \notin \{p, q\}$. Then ℓ_r does not coarse Lipschitz embed into $\ell_p \oplus \ell_q$.*

Proof. When $r > q$, the argument is based on a midpoint technique. If $r < p$, it follows immediately from Corollary 3.14. So we assume now that $1 \leq p < r < q < \infty$ and $f = (g, h) : \ell_r \rightarrow \ell_p \oplus_\infty \ell_q$ is a coarse-Lipschitz embedding. Applying the midpoint technique to the coarse Lipschitz map g and then Theorem 3.13 to the map $h \circ \varphi$ with φ of the form $\varphi(\bar{n}) = u + \tau k^{-1/r}(e_{n_1} + \cdots + e_{n_k})$, where (e_n) is the canonical basis of ℓ_r and $\tau > 0$ is large enough, leads to a contradiction. \square

We can now state and prove the main result of [63].

Theorem 3.19. *If $1 < p_1 < \cdots < p_n < \infty$ are all different from 2, then $\ell_{p_1} \oplus \cdots \oplus \ell_{p_n}$ has a unique net structure.*

Proof. We will only sketch the proof for $\ell_p \oplus \ell_q$, with $1 < p < q < \infty$ such that $2 \notin \{p, q\}$. Assume that X is a Banach space such that $X \overset{N}{\sim} \ell_p \oplus \ell_q$. The key point is again to show that X does not contain any isomorphic copy of ℓ_2 . This follows clearly from the above corollary. To conclude the proof, we need to use a few deep linear results. The cases $1 < p < q < 2$ and $2 < p < q$, were actually settled in [51] for uniform homeomorphisms. So, let us only explain the case $1 < p < 2 < q$. As in the proof of Theorem 3.17, we obtain that $X \subseteq_c L_p \oplus L_q$. Since $\ell_2 \not\subseteq X$ and $q > 2$, a theorem of Johnson [48] insures that any bounded operator from X into L_q factors through ℓ_q and therefore that $X \subseteq_c L_p \oplus \ell_q$. Then we notice that L_p and ℓ_q are totally incomparable, which means that they have no isomorphic infinite-dimensional subspaces. We can now use a theorem of Ādelštein and Wojtaszczyk [29] to obtain that $X \simeq F \oplus G$, with $F \subseteq_c L_p$ and $G \subseteq_c \ell_q$. First it follows from [91] that G is isomorphic to ℓ_q or is finite-dimensional. On the other hand, we know that $\ell_2 \not\subseteq F$, and,

by the Johnson-Odell theorem [52], F is isomorphic to ℓ_p or finite-dimensional. Summarizing, we have that X is isomorphic to ℓ_p , ℓ_q or $\ell_p \oplus \ell_q$. But we already know that ℓ_p and ℓ_q have unique net structure. Therefore, X is isomorphic to $\ell_p \oplus \ell_q$. \square

Remark. Let $1 < p < \infty$ and $p \neq 2$. It is clear that the above proof cannot work for $\ell_p \oplus \ell_2$. As was already mentioned, it is unknown whether $\ell_p \oplus \ell_2$ has a unique uniform structure. The same question is open for L_p , $1 < p < \infty$ (see Problem 1).

However, let us indicate a few other results from [63] that can be derived from Theorem 3.13. The following theorem is related to a recent result of [44] stating that, if $2 < p < \infty$, a subspace X of L_p which is not isomorphic to a subspace of $\ell_p \oplus \ell_2$ contains an isomorphic copy of $\ell_p(\ell_2)$.

Theorem 3.20. *Let $1 < p < \infty$ and $p \neq 2$. Then $\ell_p(\ell_2)$, and therefore L_p do not coarse Lipschitz embed into $\ell_p \oplus \ell_2$.*

It follows from Ribe's counterexample that reflexivity is not preserved under uniform homeomorphisms. However, the following is proved in [9].

Theorem 3.21. *Let X be a Banach space and Y a reflexive Banach space with an equivalent AUS norm. Assume that X coarse Lipschitz embeds into Y . Then X is reflexive.*

The proof has three ingredients: a result of Odell and Schlumprecht [86] asserting that Y can be renormed in such a way that $\bar{\rho}_Y \leq \bar{\rho}_{\ell_p}$, James's characterization of reflexivity and Theorem 3.13.

Remark. Note that Theorems 3.13 and 3.21 seem to take us very close to the solution of Problem 2. See also Corollary 4.6 below.

3.6. Uniform asymptotic convexity. Until very recently, there has been no corresponding result about the stability of convexity. The only thing that could be mentioned was the elementary use of the approximate midpoints principle that we already described. In a recent

article [60], Nigel Kalton made a real breakthrough in this direction. Let us first state his general result.

Theorem 3.22. *Suppose X and Y are Banach spaces and that there is a coarse Lipschitz embedding of X into Y . Then there is a constant $C > 0$ such that, for any spreading model $(e_k)_{k=1}^\infty$ of a weakly null normalized sequence in X (whose norm will be denoted $\|\cdot\|_S$), we have:*

$$\|e_1 + \cdots + e_k\|_{\widehat{\delta}_Y} \leq C \|e_1 + \cdots + e_k\|_S.$$

We will not prove this in detail. We have chosen instead to show an intermediate result whose proof contains one of the key ingredients. Let us first describe the main idea. We wish to use the approximate midpoints principle. But, unless the space is very simple or concrete (like ℓ_p spaces), the approximate midpoint set is difficult to describe. Kalton's strategy in [60] was, in order to prove the desired inequality, to define an adapted norm on an Orlicz space associated with any given weakly null sequence in X . Then, by composition, he was able to reduce the question to the study of a coarse-Lipschitz map from that space to the Orlicz space associated with the modulus of asymptotic convexity of Y . Finally, as in ℓ_p , the approximate midpoints are not so difficult to study in an Orlicz space. Before stating the result, we need some preliminary notation.

Let ϕ be an Orlicz Lipschitz function. Then, $N_\phi(1, t) = 1 + \phi(|t|)$ (for $t \in \mathbb{R}$) extends to a norm N_ϕ^2 defined on \mathbb{R}^2 . We now define inductively a norm on \mathbb{R}^j by $N_\phi^j(x_1, \dots, x_j) = N_\phi^2(N_\phi^{j-1}(x_1, \dots, x_{j-1}), x_j)$ for $j \geq 2$. These norms are compatible and define a norm Λ_ϕ on c_{00} . One can check that $1/2\|\cdot\|_\phi \leq \|\cdot\|_{\Lambda_\phi} \leq \|\cdot\|_\phi$.

We also need to introduce the following quantities:

$$\widehat{\delta}_X(t) = \inf_{x \in S_X} \sup_E \inf_{y \in S_E} \left\{ \frac{1}{2} (\|x + ty\| + \|x - ty\|) - 1 \right\}$$

where again E runs through all closed subspaces of X of finite codimension. The function $\widehat{\delta}_X(t)/t$ is increasing and so $\widehat{\delta}_X$ is equivalent to the convex function

$$\widetilde{\delta}_X(t) = \int_0^t \frac{\widehat{\delta}_X(s)}{s} ds.$$

Theorem 3.23. *Let $(\varepsilon_i)_{i=1}^\infty$ be a sequence of independent Rademacher variables on a probability space Ω . Assume that X and Y are Banach spaces and that there is a coarse Lipschitz embedding of X into Y . Then, there is a constant $c > 0$ such that, given any (x_n) weakly null normalized θ -separated sequence in X and any integer k , there exist $n_1 < \dots < n_k$ so that:*

$$c\theta\|e_1 + \dots + e_k\|_{\delta_Y} \leq \|\varepsilon_1 x_{n_1} + \dots + \varepsilon_k x_{n_k}\|_{L^1(\Omega, X)},$$

where $(e_k)_{k=1}^\infty$ is the canonical basis of c_{00} .

Proof. For $k \in \mathbb{N}$, let

$$\sigma_k = \sup \left\{ \left\| \sum_{j=1}^k \varepsilon_j x_{n_j} \right\|_{L^1(\Omega, X)}, \quad n_1 < n_2 < \dots < n_k \right\}.$$

For each k , define the Orlicz function F_k by

$$(3.1) \quad F_k(t) = \begin{cases} \sigma_k t/k, & 0 \leq t \leq 1/\sigma_k \\ t + 1/k - 1/\sigma_k, & 1/\sigma_k \leq t < \infty. \end{cases}$$

We introduce an operator $T : c_{00} \rightarrow L_1(\Omega; X)$ defined by $T(\xi) = \sum_{j=1}^\infty \xi_j \varepsilon_j \otimes x_j$. We omit the proof of the fact that, for all $\xi \in c_{00}$: $\|T\xi\| \leq 2\|\xi\|_{F_k} \leq 4\|\xi\|_{\Lambda_{F_k}}$.

Assume now that $f : X \rightarrow Y$ is a map such that $f(0) = 0$ and

$$\|x - z\| - 1 \leq \|f(x) - f(z)\| \leq K\|x - z\| + 1, \quad x, z \in X.$$

We then define $g : (c_{00}, \Lambda_{F_k}) \rightarrow L_1(\Omega; Y)$ by $g(\xi) = f \circ T\xi$. It can easily be checked that g is coarse-Lipschitz and that $\text{Lip}_\infty(g) > 0$. So we can apply the approximate midpoint principle to the map g and obtain that, for τ as large as we wish, there exist $\eta, \zeta \in c_{00}$ with $\|\eta - \zeta\|_{\Lambda_{N_k}} = 2\tau$ such that

$$g(\text{Mid}(\eta, \zeta, 1/k)) \subset \text{Mid}(g(\eta), g(\zeta), 2/k).$$

Let $m \in \mathbb{N}$ so that $\eta, \zeta \in \text{span}\{e_1, \dots, e_{m-1}\}$. It follows from the definition of Λ_{N_k} that, for $j \geq m$: $\xi + \tau\sigma_k^{-1}e_j \in \text{Mid}(\eta, \zeta, 1/k)$, where $\xi = (\eta + \zeta)/2$. Thus, the functions

$$h_j = f \circ \left(\sum_{i=1}^{m-1} \xi_i \varepsilon_i \otimes x_i + \tau\sigma_k^{-1} \varepsilon_j \otimes x_j \right), \quad j \geq m$$

all belong to $\text{Mid}(g(\eta), g(\zeta), 2/k)$. Since the ε_i 's are independent so are the functions

$$h'_j = f\left(\sum_{i=1}^{m-1} \xi_i \varepsilon_i \otimes x_i + \tau \sigma_k^{-1} \varepsilon_m \otimes x_j\right), \quad j \geq m.$$

Therefore, for all $j \geq m$:

$$\|g(\eta) - h'_j\| + \|g(\zeta) - h'_j\| - \|g(\eta) - g(\zeta)\| \leq 2k^{-1} \|g(\eta) - g(\zeta)\|.$$

We shall now use without proof the following simple property of $N = N_{\delta_Y}^2$: for any bounded t -separated sequence in Y and any $z \in Y$,

$$(3.2) \quad \liminf_{n \rightarrow \infty} (\|y - y_n\| + \|z - y_n\|) \geq N(\|y - z\|, t).$$

Note that, for any $\omega \in \Omega$, we have

$$\|h'_i(\omega) - h'_j(\omega)\| \geq \theta \tau \sigma_k^{-1} - 1, \quad i > j \geq m.$$

Hence, using (3.2), integrating and using Jensen's inequality, we get

$$\liminf_{j \rightarrow \infty} (\|g(\eta) - h'_j\| + \|g(\zeta) - h'_j\|) \geq N(\|g(\eta) - g(\zeta)\|, \theta \tau \sigma_k^{-1} - 1).$$

Now $\|g(\eta) - g(\zeta)\| \leq 8K\tau + 1$ and $N(t, 1) - t$ is a decreasing function, so

$$\begin{aligned} N_Y(8K\tau + 1, \theta \tau \sigma_k^{-1} - 1) - (8K\tau + 1) &\leq \frac{2}{k} \|g(\eta) - g(\zeta)\| \\ &\leq 2(8K\tau + 1)k^{-1}. \end{aligned}$$

Multiplying by $(8K\tau + 1)^{-1}$ and letting τ tend to $+\infty$, we obtain that

$$\tilde{\delta}\left(\frac{\theta}{16K\sigma_k}\right) \leq \frac{2}{k} \quad \text{and therefore} \quad \tilde{\delta}\left(\frac{\theta}{32K\sigma_k}\right) \leq \frac{1}{k}$$

or

$$\|e_1 + \cdots + e_k\|_{\tilde{\delta}_Y} \leq 32K\theta^{-1}\sigma_k. \quad \square$$

We will end this section by stating two theorems proved by Kalton in [60]. Their proofs use, among many other ideas, the results we just explained on the stability of asymptotic uniform convexity under coarse-Lipschitz embeddings.

Theorem 3.24. [?] *Suppose $1 < p < \infty$. Then:*

- (i) *If X is a Banach space which can be coarse Lipschitz embedded in ℓ_p , then X is linearly isomorphic to a closed subspace of ℓ_p .*
- (ii) *If X is a Banach space which is net equivalent to a quotient of ℓ_p , then X is linearly isomorphic to a quotient of ℓ_p .*
- (iii) *If X can be coarse Lipschitz embedded into a quotient of ℓ_p , then X is linearly isomorphic to a subspace of a quotient of ℓ_p .*

Remark. On the other hand, Kalton constructed in [61] two subspaces (respectively, quotients) of ℓ_p ($1 < p \neq 2 < \infty$) which are uniformly homeomorphic but not linearly isomorphic.

Problem 4. It is not known whether a Banach space which is net equivalent to a subspace (respectively a quotient) of c_0 is linearly isomorphic to a subspace (respectively a quotient) of c_0 .

As we have already seen, a Banach space Lipschitz-isomorphic to a subspace of c_0 is linearly isomorphic to a subspace of c_0 [38]. It is not known if the class of Banach spaces linearly isomorphic to a quotient of c_0 is stable under Lipschitz-isomorphisms. However, this question was almost solved by Dutrieux who proved in [26] that, if a Banach space is Lipschitz-isomorphic to a quotient of c_0 and has a dual with the approximation property, then it is linearly isomorphic to a quotient of c_0 .

Finally, let us point out one last striking consequence of more general results from [60].

Theorem 3.25. [?] *Suppose $1 < p, r < \infty$ are such that $p < \min(r, 2)$ or $p > \max(r, 2)$. Then the space $(\sum_{n=1}^{\infty} \ell_r^n)_{\ell_p}$ has a unique net structure.*

4. Embeddings of special graphs into Banach spaces. In this section we will study special metric graphs or trees that are of particular importance for the subject. More precisely, we will study the Banach spaces in which they embed. This will allow us to characterize some linear classes of Banach spaces by a purely metric condition of the following type: given a metric space M (generally a graph), what are the Banach spaces X so that $M \xrightarrow{\text{Lip}} X$. Or, given a family \mathcal{M} of metric

spaces, what are the Banach spaces X for which there is a constant $C \geq 1$ so that, for all M in \mathcal{M} , $M \xrightarrow{C} X$ (i.e., M Lipschitz embeds into X with distortion at most C).

Most of the time, these linear classes were already known to be stable under Lipschitz or coarse-Lipschitz embeddings, when such characterizations were proved. However, we will show one situation (see Corollary 4.6) where this process yields new results about such stabilities.

The section will be organized by the nature of the linear properties that can be characterized by such embedding conditions.

4.1. Embeddings of special metrics spaces and local properties of Banach spaces.

We already know (see Theorem 3.3) that local properties of Banach spaces are preserved under coarse Lipschitz embeddings. This theorem gave birth to the ‘‘Ribe program’’ which aims at looking for metric invariants that characterize local properties of Banach spaces. Recently, a detailed description of the Ribe program was given by Naor in [81]. The first occurrence of the Ribe program was Bourgain’s metric characterization of superreflexivity given in [17]. The metric invariant discovered by Bourgain is the collection of the hyperbolic dyadic trees of arbitrarily large height N . We denote $\Delta_0 = \{\emptyset\}$ as the root of the tree. Let $\Omega_i = \{-1, 1\}^i$, $\Delta_N = \bigcup_{i=0}^N \Omega_i$ and $\Delta_\infty = \bigcup_{i=0}^\infty \Omega_i$. Then we equip Δ_∞ , and by restriction every Δ_N , with the hyperbolic distance ρ , which is defined as follows. Let s and s' be two elements of Δ_∞ , and let $u \in \Delta_\infty$ be their greatest common ancestor. We set

$$\rho(s, s') = |s| + |s'| - 2|u| = \rho(s, u) + \rho(s', u).$$

Bourgain’s characterization is the following:

Theorem 4.1. [17]. *Let X be a Banach space. Then X is not superreflexive if and only if there exists a constant $C \geq 1$ such that, for all $N \in \mathbb{N}$, $(\Delta_N, \rho) \xrightarrow{C} X$.*

Remark. It has been proved by Baudier in [8] that this is also equivalent to the metric embeddability of the infinite hyperbolic dyadic

tree (Δ_∞, ρ) . It should also be noted that, in [8, 17], the embedding constants are bounded above by a universal constant.

We also wish to mention that Johnson and Schechtman [54] recently characterized the super-reflexivity through the non embeddability of other graphs such as the “diamond graphs” or the Laakso graphs. We will only give an intuitive description of the diamond graphs. D_0 is made of two connected vertices (therefore, at distance 1), that we shall call T (top) and B (bottom). D_1 is a diamond, therefore made of four vertices T , B , L (left) and R (right) and four edges: $[B, L]$, $[L, T]$, $[T, R]$ and $[R, B]$. Assume D_N is constructed; then D_{N+1} is obtained by replacing each edge of D_N by a diamond D_1 . The distance on D_{N+1} is the path metric of this new discrete graph. The graph distance on a diamond D_N will be denoted by d . The result is the following.

Theorem 4.2. [54]. *Let X be a Banach space. Then X is not super-reflexive if and only if there is a constant $C \geq 1$ such that, for all $N \in \mathbb{N}$, $(D_N, d) \xrightarrow{C} X$.*

The metric characterization of the linear type of a Banach space has been initiated by Enflo in [31] and continued by Bourgain, Milman and Wolfson in [18]. Let us first describe a concrete result from [18]. For $1 \leq p \leq 2$ and $n \in \mathbb{N}$, H_p^n denotes the set $\{0, 1\}^n$ equipped with the metric induced by the ℓ_p norm. The metric space H_1^n is called the Hamming cube. One of their results is the following.

Theorem 4.3. [18]. *Let X be a Banach space and $1 \leq p \leq 2$. Define p_X to be the supremum of all r 's such that X is of linear type r . Then, the following assertions are equivalent.*

- (i) $p_X \leq p$.
- (ii) *There is a constant $C \geq 1$ such that, for all $n \in \mathbb{N}$, $H_p^n \xrightarrow{C} X$.*

In particular, X is of trivial type if and only if $H_1^n \xrightarrow{C} X$, for all $n \in \mathbb{N}$ and for some universal constant $C \geq 1$.

The fundamental problem of defining a notion of type for metric spaces is behind this result. Of course we expect such a notion to

coincide with the linear type for Banach spaces and to be stable under reasonable non linear embeddings. This program was achieved with the successive definitions of the Enflo type [31], the Bourgain-Milman-Wolfson type [18] and finally the scaled Enflo type introduced by Mendel and Naor in [77]. An even more difficult task was to define the right notion of metric cotype. This was achieved by Mendel and Naor in [78]. We will not address this subject in this survey, but we advise the interested reader to study these fundamental papers.

Let us also describe a simpler metric characterization of the Banach spaces without (linear) cotype. First, let us recall that a metric space (M, d) is called *locally finite* if all its balls of finite radius are finite. It is of *bounded geometry* if, for any $r > 0$, there exists $C(r) \in \mathbb{N}$ such that the cardinal of any ball of radius r is less than $C(r)$. We will now construct a particular metric space with bounded geometry. For $k, n \in \mathbb{N}$, denote

$$M_{n,k} = knB_{\ell_\infty^n} \cap n\mathbb{Z}^n.$$

Let us enumerate the $M_{n,k}$'s: M_1, \dots, M_i, \dots with $M_i = M_{n_i, k_i}$ in such a way that $\text{diam}(M_i)$ is non decreasing. Note that $\lim_i \text{diam}(M_i) = +\infty$. Then, let M_0 be the disjoint union of the M_i 's ($i \geq 1$), and define on M_0 the following distance:

If $x, y \in M_i$, $d(x, y) = \|x - y\|_\infty$, where $\|\cdot\|_\infty$ is the $\ell_\infty^{n_i}$ norm.

If $x \in M_i$ and $y \in M_j$, with $i < j$, set $d(x, y) = F(j)$, where F is built so that F is increasing and, for all j for all $i \leq j$,

$$F(j) \geq \frac{1}{2} \text{diam}(M_j).$$

Note that $\lim_i F(i) = +\infty$. We leave it to the reader to check that (M_0, d) is a metric space with bounded geometry. We can now state the following.

Theorem 4.4. *Let X be a Banach space. The following assertions are equivalent.*

- (i) X has a trivial cotype.
- (ii) X contains uniformly and linearly the ℓ_∞^n 's.
- (iii) There exists $C \geq 1$ such that, for every locally finite metric space M , $M \xrightarrow{C} X$.

- (iv) *There exists $C \geq 1$ such that, for every metric space with bounded geometry M , $M \xrightarrow{C} X$.*
- (v) *There exists $C \geq 1$ such that $M_0 \xrightarrow{C} X$.*
- (vi) *There exists $C \geq 1$ such that, for every finite metric space M , $M \xrightarrow{C} X$.*

Proof. The equivalence between (i) and (ii) is part of classical results by Maurey and Pisier [74].

(ii) \Rightarrow (iii) is due to Baudier and the second author [10].

(iii) \Rightarrow (iv) and (iv) \Rightarrow (v) are trivial.

For any $k, n \in \mathbb{N}$, the space M_0 contains the space $M_{n,k} = knB_{\ell_\infty^n} \cap n\mathbb{Z}^n$ which is isometric to the $1/k$ -net of $B_{\ell_\infty^n} : B_{\ell_\infty^n} \cap 1/k\mathbb{Z}^n$. But, after rescaling, any finite metric space is isometric to a subset of $B_{\ell_\infty^n}$, for some $n \in \mathbb{N}$. Thus, for any finite metric space M and any $\varepsilon > 0$, there exist $k, n \in \mathbb{N}$ so that M is $(1 + \varepsilon)$ -equivalent to a subset of $M_{k,n}$. The implication (v) \Rightarrow (vi) is now clear.

The proof of (vi) \Rightarrow (ii) relies on an argument due to Schechtman [98]. So, assume that (vi) is satisfied, and let us fix $n \in \mathbb{N}$. Then, for any $k \in \mathbb{N}$, there exists a map $f_k : (1/k\mathbb{Z}^n \cap B_{\ell_\infty^n}, \|\cdot\|_\infty) \rightarrow X$ such that $f_k(0) = 0$ and for all $x, y \in 1/k\mathbb{Z}^n \cap B_{\ell_\infty^n}$,

$$\|x - y\|_\infty \leq \|f_k(x) - f_k(y)\| \leq K\|x - y\|_\infty.$$

Then, we can define a map $\lambda_k : B_{\ell_\infty^n} \rightarrow 1/k\mathbb{Z}^n \cap B_{\ell_\infty^n}$ such that, for all $x \in B_{\ell_\infty^n} : \|\lambda_k(x) - x\|_\infty = d(x, 1/k\mathbb{Z}^n \cap B_{\ell_\infty^n})$. We now set $\varphi_k = f_k \circ \lambda_k$. Let \mathcal{U} be a non trivial ultrafilter. We define $\varphi : B_{\ell_\infty^n} \rightarrow X_{\mathcal{U}} \subseteq X_{\mathcal{U}}^{**}$ by $\varphi(x) = (\varphi_k(x))_{\mathcal{U}}$. It is easy to check that φ is a Lipschitz embedding. Then it follows from results by Heinrich and Mankiewicz on weak*-Gâteaux differentiability of Lipschitz maps [45] that ℓ_∞^n is K -isomorphic to a linear subspace of $X_{\mathcal{U}}^{**}$. Finally, using the local reflexivity principle and properties of the ultra-product, we get that ℓ_∞^n is $(K + 1)$ -isomorphic to a linear subspace of X . \square

Remark. The following generalization of the implication (ii) \Rightarrow (iii) is proved in [88]: if A is locally finite metric space whose finite subsets uniformly bilipschitz (respectively, coarsely) embed into a Banach space X , then A bilipschitz (respectively, coarsely) embeds into X . Let us

also mention that other characterizations of classes of Banach spaces in terms of bilipschitz embeddability of graphs of bounded degree were also found for instance in [89].

4.2. Embeddings of special graphs and asymptotic structure of Banach spaces.

We will start this section by considering the countably branching hyperbolic trees. For a positive integer N , $T_N = \bigcup_{i=0}^N \mathbb{N}^i$, where $\mathbb{N}^0 := \{\emptyset\}$. Then $T_\infty = \bigcup_{N=1}^\infty T_N$ is the set of all finite sequences of positive integers. The hyperbolic distance ρ is defined on T_∞ as follows. Let s and s' be two elements of T_∞ , and let $u \in T_\infty$ be their greatest common ancestor. We set

$$\rho(s, s') = |s| + |s'| - 2|u| = \rho(s, u) + \rho(s', u).$$

The following result, that appeared in [9], is an asymptotic analogue of Bourgain's characterization of super-reflexivity given in Theorem 4.1 above.

Theorem 4.5. *Let X be a reflexive Banach space. The following assertions are equivalent.*

- (i) *There exists $C \geq 1$ such that $T_\infty \xrightarrow{C} X$.*
- (ii) *There exists $C \geq 1$ such that for any N in \mathbb{N} , $T_N \xrightarrow{C} X$.*
- (iii) *X does not admit any equivalent asymptotically uniformly smooth norm or X does not admit any equivalent asymptotically uniformly convex norm.*

We will only mention one application of this result.

Corollary 4.6. *The class of all reflexive Banach spaces that admit both an equivalent AUS norm and an equivalent AUC norm is stable under coarse Lipschitz embeddings.*

Proof. Assume that X coarse Lipschitz embeds in a space Y which is reflexive, AUS renormable and AUC renormable. First, it follows from Theorem 3.21 that X is reflexive. Now the conclusion is easily derived from Theorem 4.5. \square

Note that this class coincides with the class of reflexive spaces X such that the Szlenk indices of X and X^* are both equal to the first infinite ordinal ω (see [39]).

Problem 5. We do not know if the class of all Banach spaces that are both AUS renormable and AUC renormable is stable under coarse Lipschitz embeddings, net equivalences or uniform homeomorphisms.

Problem 6. We now present a variant of Problem 2. As we already indicated, we do not know if the class of reflexive and AUS renormable Banach spaces is stable under coarse Lipschitz embeddings. The important results by Kalton and Randrianarivony on the stability of the asymptotic uniform smoothness are based on the use of particular metric graphs, namely the graphs $G_k(\mathbb{N})$ equipped with the distance: for all $\bar{n}, \bar{m} \in G_k(\mathbb{N})$,

$$d(\bar{n}, \bar{m}) = |\{j, n_j \neq m_j\}|.$$

It seems interesting to try to characterize the Banach spaces X such that there exists a constant $C \geq 1$ for which $G_k(\mathbb{N}) \xrightarrow{C} X$, for all $k \in \mathbb{N}$. In particular, one may ask whether a reflexive Banach space which is not AUS renormable always contains the $G_k(\mathbb{N})$'s with uniform distortion (the converse implication is a consequence of Kalton and Randrianarivony's work). A positive answer would solve Problem 2.

4.3. Interlaced Kalton's graphs. Very little is known about the coarse embeddings of metric spaces into Banach spaces and about coarse embeddings between Banach spaces (see Definition 3.1 for coarse embeddings). For quite some time it was not even known if a reflexive Banach space could be universal for separable metric spaces and coarse embeddings. This was solved negatively by Kalton in [56] who showed the following.

Theorem 4.7. [56]. *Let X be a separable Banach space. Assume that c_0 coarsely embeds into X . Then one of the iterated duals of X has to be non separable. In particular, X cannot be reflexive.*

The idea of the proof is to consider a new graph metric δ on $G_k(\mathbb{M})$, for \mathbb{M} infinite subset of \mathbb{N} . We will say that $\bar{n} \neq \bar{m} \in G_k(\mathbb{M})$ are

adjacent (or $\delta(\bar{n}, \bar{m}) = 1$) if they interlace or, more precisely, if

$$m_1 \leq n_1 \leq \dots \leq m_k \leq n_k \text{ or } n_1 \leq m_1 \leq \dots \leq n_k \leq m_k.$$

For simplicity, we will only show that X cannot be reflexive. So, let us assume that X is a reflexive Banach space and fix a non principal ultrafilter \mathcal{U} on \mathbb{N} . For a bounded function $f : G_k(\mathbb{N}) \rightarrow X$, we define $\partial f : G_{k-1}(\mathbb{N}) \rightarrow X$ by, for all $\bar{n} \in G_{k-1}(\mathbb{N})$,

$$\partial f(\bar{n}) = w - \lim_{n_k \in \mathcal{U}} f(n_1, \dots, n_{k-1}, n_k).$$

Note that, for $1 \leq i \leq k$, $\partial^i f$ is a bounded map from $G_{k-i}(\mathbb{N})$ into X and that $\partial^k f$ is an element of X . Let us first state without proof a series of basic lemmas about the operation ∂ .

Lemma 4.8. *Let $h : G_k(\mathbb{N}) \rightarrow \mathbb{R}$ be a bounded map and $\varepsilon > 0$. Then there is an infinite subset \mathbb{M} of \mathbb{N} such that, for all $\bar{n} \in G_k(\mathbb{M})$,*

$$|h(\bar{n}) - \partial^k h| < \varepsilon.$$

Lemma 4.9. *Let $f : G_k(\mathbb{N}) \rightarrow X$ and $g : G_k(\mathbb{M}) \rightarrow X^*$ be two bounded maps. Define $f \otimes g : G_{2k}(\mathbb{N}) \rightarrow \mathbb{R}$ by*

$$(f \otimes g)(n_1, \dots, n_{2k}) = \langle f(n_2, n_4, \dots, n_{2k}), g(n_1, \dots, n_{2k-1}) \rangle.$$

Then $\partial^2(f \otimes g) = \partial f \otimes \partial g$.

Lemma 4.10. *Let $f : G_k(\mathbb{N}) \rightarrow X$ be a bounded map and $\varepsilon > 0$. Then there is an infinite subset \mathbb{M} of \mathbb{N} such that, for all $\bar{n} \in G_k(\mathbb{M})$,*

$$\|f(\bar{n})\| \leq \|\partial^k f\| + \omega_f(1) + \varepsilon,$$

where ω_f is the modulus of continuity of f .

Lemma 4.11. *Let $\varepsilon > 0$, X a separable reflexive Banach space and I an uncountable set. Assume that, for each $i \in I$, $f_i : G_k(\mathbb{N}) \rightarrow X$ is a bounded map. Then there exist $i \neq j \in I$ and an infinite subset \mathbb{M} of \mathbb{N} such that, for all $\bar{n} \in G_k(\mathbb{M})$,*

$$\|f_i(\bar{n}) - f_j(\bar{n})\| \leq \omega_{f_i}(1) + \omega_{f_j}(1) + \varepsilon.$$

We are now ready for the proof of Theorem 4.7. As we will see, it relies on the fact that c_0 contains uncountably many isometric copies

of the $G_k(\mathbb{N})$'s with too many points far away from each other (which will be in contradiction with Lemma 4.11).

Proof of Theorem 4.7. Assume X is reflexive, and let $h : c_0 \rightarrow X$ be a map which is bounded on bounded subsets of c_0 . Let $(e_k)_{k=1}^\infty$ be the canonical basis of c_0 . For an infinite subset A of \mathbb{N} , we now define, for all $n \in \mathbb{N}$,

$$s_A(n) = \sum_{k \leq n, k \in A} e_k$$

and, for all $\bar{n} = (n_1, \dots, n_k) \in G_k(\mathbb{N})$,

$$f_A(\bar{n}) = \sum_{i=1}^k s_A(n_i).$$

Then the $h \circ f_A$'s form an uncountable family of bounded maps from $G_k(\mathbb{N})$ to X . It therefore follows from Lemma 4.11 that there are two distinct infinite subsets A and B of \mathbb{N} and another infinite subset \mathbb{M} of \mathbb{N} so that: for all $\bar{n} \in G_k(\mathbb{M})$

$$\begin{aligned} \|h \circ f_A(\bar{n}) - h \circ f_B(\bar{n})\| &\leq \omega_{h \circ f_A}(1) + \omega_{h \circ f_B}(1) + 1 \\ &\leq 2\omega_h(1) + 1. \end{aligned}$$

But, since $A \neq B$, there is $\bar{n} \in G_k(\mathbb{M})$ with $\|f_A(\bar{n}) - f_B(\bar{n})\| = k$. By taking arbitrarily large values of k , we deduce that h cannot be a coarse embedding. \square

Remarks.

- (1) Similarly, one can show that h cannot be a uniform embedding, by composing h with the maps tf_A and letting t tend to zero.
- (2) It is now easy to adapt this proof in order to obtain the stronger result stated in Theorem 4.7. Indeed, one just has to change the definition of the operator ∂ as follows. If $f : G_k(\mathbb{N}) \rightarrow X$ is bounded, define $\partial f : G_{k-1}(\mathbb{N}) \rightarrow X^{**}$ by, for all $\bar{n} \in G_{k-1}(\mathbb{N})$,

$$\partial f(\bar{n}) = w^* - \lim_{n_k \in \mathcal{U}} f(n_1, \dots, n_{k-1}, n_k).$$

We leave it to the reader to rewrite the argument.

- (3) On the other hand, Kalton proved in [55] that c_0 embeds uniformly and coarsely in a Banach space X with the Schur

property. Note that X does not contain any subspace linearly isomorphic to c_0 .

- (4) We will see in the next section that Kalton recently used a similar operation ∂ and the same graph distance on $G_k(\omega_1)$, where ω_1 is the first uncountable ordinal (see [58]) in order to study uniform embeddings into ℓ_∞ .

Problem 7. In view of this result, the metric graphs $(G_k(\mathbb{N}), \delta)$ are clearly of special importance. It is a natural question to characterize the Banach spaces containing the spaces $(G_k(\mathbb{N}), \delta)$ with uniformly bounded distortion.

In [56], Kalton pushed the idea behind the proof of Theorem 4.7 much further and introduced the following abstract notions in order to study the coarse or uniform embeddings into reflexive Banach spaces.

Let (M, d) be a metric space, $\varepsilon > 0$ and $\eta \geq 0$. We say that M has *property* $\mathcal{Q}(\varepsilon, \eta)$ if for every $k \in \mathbb{N}$, and for every map $f : (G_k(\mathbb{N}), \delta) \rightarrow (M, d)$ with $\omega_f(1) \leq \eta$ there exists an infinite subset \mathbb{M} of \mathbb{N} such that:

$$d(f(\sigma), f(\tau)) \leq \varepsilon \quad \sigma < \tau, \quad \sigma, \tau \in G_k(\mathbb{M}).$$

Then $\Delta_M(\varepsilon)$ is the supremum of all $\eta \geq 0$ so that M has *property* $\mathcal{Q}(\varepsilon, \eta)$ and Kalton proves the following general statement.

Theorem 4.12. *Let M be a metric space and X be a reflexive Banach space.*

- (i) *If M embeds uniformly into X , then $\Delta_M(\varepsilon) > 0$, for all $\varepsilon > 0$.*
- (ii) *If M embeds coarsely into X , then $\lim_{\varepsilon \rightarrow +\infty} \Delta_M(\varepsilon) = +\infty$.*

Let us now turn to the case when our metric space is a Banach space that we shall denote E . Then it is easy to see that the function Δ_E is linear. We denote \mathcal{Q}_E the constant such that, for all $\varepsilon > 0$, $\Delta_E(\varepsilon) = \mathcal{Q}_E \varepsilon$. Finally, we say that E has the *\mathcal{Q} -property* if $\mathcal{Q}_E > 0$. It follows from Lemma 4.10 that a reflexive Banach space has the *\mathcal{Q} -property*. Thus, we have:

Corollary 4.13. *If a Banach space E fails the \mathcal{Q} -property, then E does not coarsely embed into a reflexive Banach space and B_E does not uniformly embed into a reflexive Banach space.*

The fact that c_0 fails the \mathcal{Q} -property follows from Theorem 4.7, but it is actually an ingredient of its proof. Then Kalton continues his study of the links between reflexivity and the \mathcal{Q} -property. Let us mention without proof a few of the many interesting results obtained in [56].

- (1) A non reflexive Banach space with the alternating Banach-Saks property (see the definition in [56]), in particular with a non trivial type, fails the \mathcal{Q} -property.
- (2) The James space J and its dual fail the \mathcal{Q} -property.
- (3) However, there exists a quasi-reflexive but non reflexive Banach space with the \mathcal{Q} -property.

Problem 8. Is there a converse to Corollary 4.13? More precisely, if E is a separable Banach space with the \mathcal{Q} -property, does B_E uniformly embed into a reflexive Banach space or does E coarsely embed into a reflexive Banach space? The answer is unknown for the space constructed in the above statement (3).

5. Nonseparable spaces. We collect here a few recent results obtained by Nigel Kalton on nonseparable Banach spaces together with some related open problems. All the results presented in this section are taken from Kalton's paper [58]. They mainly concern embeddings of nonseparable Banach spaces into ℓ_∞ . We start with a positive result.

Theorem 5.1. *If X has an unconditional basis and is of density character at most c (the cardinal of the continuum), then it is Lipschitz embeddable into ℓ_∞ .*

Sketch of the main ideas in the proof. Assume that the unconditionality constant of the basis is 1 and that it is indexed by the set \mathbb{R} of real numbers. Denote by $(e_t^*)_{t \in \mathbb{R}}$ the biorthogonal functionals of the basis. If $x \in X$, we write $x(t) = e_t^*(x)$. Suppose that $a, b, c \in \mathbb{Q}^n$. We write typically, $a = (a_1, a_2, \dots, a_n)$ and denote by $-a = (-a_1, -a_2, \dots, -a_n)$.

Then define a subset $U(a, b, c) \subset \mathbb{R}^n$ by $(t_1, t_2, \dots, t_n) \in U(a, b, c)$ if $b_j < t_j < c_j$ for $j = 1, 2, \dots, n$, $t_1 < t_2 < \dots, t_n$ and

$$\left\| \sum_{j=1}^n a_j e_{t_j}^* \right\|_{X^*} \leq 1.$$

For $t \in \mathbb{R}$, write $t_+ = \max(t, 0)$, and define $f_{(a,b,c)} : X \rightarrow \mathbb{R}$ by $f_{(a,b,c)}$ identically 0 if $U(a, b, c)$ is empty and, otherwise,

$$f_{(a,b,c)}(x) = \sup \left\{ \sum_{j=1}^n (a_j x(t_j))_+, (t_1, t_2, \dots, t_n) \in U(a, b, c) \right\}.$$

Finally, define the map

$$F(x) = (f_{(a,b,c)}(x))_{(a,b,c) \in \bigcup_n Q^n}$$

It can then be shown that F is a Lipschitz embedding of X into ℓ_∞ . \square

Problem 9. Let X be reflexive of density $\leq c$. Is X Lipschitz embeddable in ℓ_∞ ?

We now proceed with other spaces of density $\leq c$. Let I be a set of cardinality c . It is easy to show, using almost disjoint families, that the space $c_0(I)$ is isometric to a subspace of ℓ_∞/c_0 , and it follows that there is no linear continuous injective map from ℓ_∞/c_0 into ℓ_∞ . But, by Theorem 5.1 above, $c_0(I)$ Lipschitz embeds into ℓ_∞ . This was shown much earlier [2] using the space JL_∞ , and it can also be seen by applying [24, Theorem VI.8.9] to any separable compact space K with weight c and with some finite derivative empty. Hence, the linear argument does not extend to the non-linear case. However, Kalton showed:

Theorem 5.2. $C[0, \omega_1]$ or ℓ_∞/c_0 cannot be uniformly embedded into ℓ_∞ .

Before discussing the main ideas in the proof of this result, we need some preparation. For $n \geq 0$, let $\Omega_n = \Omega_1^{[n]}$ be the collection of all n -subsets of $\Omega_1 = [1, \omega_1)$. For $n = 0$, $\Omega_0 = \{\emptyset\}$. We write a typical element of Ω_n in the form $\alpha = \{\alpha_1, \dots, \alpha_n\}$, where $\alpha_1 < \alpha_2 < \dots < \alpha_n$. If $n \geq 1$, and $A \subset \Omega_n$, we define $\partial A \subset \omega_1^{[n-1]}$ by $\{\alpha_1, \dots, \alpha_{n-1}\} \in \partial A$ if and only if $\{\beta : \{\alpha_1, \dots, \alpha_{n-1}, \beta\} \in A\}$ is uncountable. If $n = 1$, this means that $\emptyset \in \partial A$ if and only if A is uncountable.

We will say that $A \subset \Omega_n$ is *large* if $\emptyset \in \partial^n A$. Otherwise, A is *small*. We will say that $A \subset \Omega_n$ is *very large* if its complement is small. Then one can show the following Ramsey type result.

Lemma 5.3. *If A is a very large subset in Ω_n , then there is an uncountable set $\Theta \subset \Omega_1$ such that $\Theta^{[n]} \subset A$.*

We will now make Ω_n into a graph by declaring $\alpha \neq \beta$ to be adjacent if they interlace, namely, if

$$\alpha_1 \leq \beta_1 \leq \cdots \leq \alpha_n \leq \beta_n \quad \text{or} \quad \beta_1 \leq \alpha_1 \leq \cdots \leq \beta_n \leq \alpha_n,$$

and we define d to be the least path metric on Ω_n , which then becomes a metric space.

We write $\alpha < \beta$ if $\alpha_1 < \cdots < \alpha_n < \beta_1 < \cdots < \beta_n$. If $\alpha < \beta$, then $d(\alpha, \beta) = n$ so Ω_n has diameter n .

The next lemma relies on basic properties of the ordered set Ω_1 .

Lemma 5.4.

- (i) *If A and B are large sets in Ω_n , then there exist $\alpha \in A$ and $\beta \in B$ so that α and β interlace.*
- (ii) *If f is a Lipschitz map from Ω_n into \mathbb{R} , with Lipschitz constant L , then there is $\xi \in \mathbb{R}$ so that $\{\alpha, |f(\alpha) - \xi| > L/2\}$ is small.*

It yields:

Proposition 5.5. *If f is a Lipschitz map from $(\Omega_n, d) \rightarrow \ell_\infty$ with Lipschitz constant L , then there are $\xi \in \ell_\infty$ and an uncountable subset Θ of Ω_1 so that*

$$\|f(\alpha) - \xi\| \leq L/2, \quad \alpha \in \Theta^{[n]}.$$

Sketch of the proof of Theorem 5.2. For the case of $C[0, \omega_1]$, let $(x_\mu)_{\mu \leq \omega_1}$ be defined by $x_\mu = \chi_{[0, \mu]}$. Assume that $X = C[0, \omega_1]$ uniformly homeomorphically embeds into ℓ_∞ , and let $f : B_X \rightarrow \ell_\infty$ be a uniformly homeomorphic embedding. Under the notation above, for each n , consider the map $f_n : \Omega_n \rightarrow \ell_\infty$ given by

$$f_n(\alpha) = f\left(\frac{1}{n} \sum_{j=1}^n x_{\alpha_j}\right)$$

If α and β interlace, then by a telescopic argument,

$$\left\| \frac{1}{n} \sum_{j=1}^n (x_{\beta_j} - x_{\alpha_j}) \right\| \leq \frac{2}{n},$$

and from the definition of the distance in Ω_n , we get that f_n has Lipschitz constant $\psi_f(2/n)$, where ψ_f is the modulus of uniform continuity of f .

By Proposition 5.5, we may pick an uncountable subset Θ_n of Ω_1 so that

$$\|f_n(\alpha) - f_n(\beta)\| \leq \psi_f\left(\frac{2}{n}\right), \quad \alpha, \beta \in \Theta_n.$$

Hence,

$$\left\| \frac{2}{n} \sum_{j=1}^n x_{\beta_j} - \frac{2}{n} \sum_{j=1}^n x_{\alpha_j} \right\| \leq \psi_g\left(\psi_f\left(\frac{2}{n}\right)\right), \quad \alpha, \beta \in \Theta_n.$$

Now pick $\alpha_1 < \alpha_2 < \dots < \alpha_n \in \Theta_n$. If $\nu > \mu > \alpha_n$, we can find β_1, \dots, β_n in Θ_n so that $\beta_n > \beta_{n-1} > \dots > \beta_1 > \nu$. Then

$$\|x_\nu - x_\mu\| \leq \left\| \frac{1}{n} \sum_{j=1}^n x_{\beta_j} - \frac{1}{n} \sum_{j=1}^n x_{\alpha_j} \right\| \leq \psi_g\left(\psi_f\left(\frac{2}{n}\right)\right).$$

Thus,

$$\theta(\mu) := \sup_{\sigma > \mu} \|x_\sigma - x_\mu\| \leq \psi_g\left(\psi_f\left(\frac{2}{n}\right)\right), \quad \mu > \alpha_n.$$

Applying this for every n , since $\lim_{n \rightarrow \infty} \psi_g(\psi_f(2/n)) = 0$, we get $\theta(\mu) = 0$ eventually, which is not true.

For ℓ_∞/c_0 , Theorem 5.2 follows from the case of $C[0, \omega_1]$ and from the result that $C[0, \omega_1]$ is linearly isometric to a subspace of ℓ_∞/c_0 [90]. \square

Note that Theorem 5.2 implies that there is no quasi-additive Lipschitz projection from ℓ_∞ onto c_0 (see [13]). As another application of Theorem 5.2, Kalton obtains the following fundamental example.

Theorem 5.6. *There is a (nonseparable) Banach space Z that is not a uniform retract of its second dual.*

Before starting discussion of the ideas in the proof, let us include the following useful lemma of independent interest.

Lemma 5.7. *Let X be a Banach space, and let $Q : Y \rightarrow X$ be a quotient mapping. In order that there is a uniformly continuous selection $f : B_X \rightarrow Y$ of the quotient mapping Q , it is sufficient that for some $0 < \lambda < 1$ there is a uniformly continuous map $\phi : S_X \rightarrow Y$ with $\|Q(\phi(x)) - x\| \leq \lambda$ for $x \in S_X$.*

Proof. We extend ϕ to B_X to be positively homogeneous and ϕ remains uniformly continuous. Define $g(x) = x - Q(\phi(x))$, so that g is also positively homogeneous. Then $\|g(x)\| \leq \lambda\|x\|$, and so $\|g^n(x)\| \leq \lambda^n\|x\|^n$ for $x \in B_X$. Let $g^0(x) = x$. Let

$$f(x) = \sum_{n=0}^{\infty} \phi(g^n(x)).$$

The series converges uniformly in $x \in B_X$, and so f is uniformly continuous. Furthermore,

$$Qf(x) = \sum_{n=0}^{\infty} (g^n(x) - g^{n+1}(x)) = x. \quad \square$$

Sketch of the proof of Theorem 5.6. The space Z that we shall consider was constructed by Benyamini in [12]. Consider the quotient map $Q : \ell_\infty \rightarrow \ell_\infty/c_0$. For each n , pick a maximal set D_n in the interior of B_{ℓ_∞/c_0} so that $\|x - x'\| \geq 1/n$ for $x, x' \in D_n$ and $x \neq x'$. Then, for each n , define a map $h_n : D_n \rightarrow B_{\ell_\infty}$ with $Qh_n(x) = x$ for $x \in D_n$, and denote by Y_n the space ℓ_∞ with the equivalent norm

$$\|y\|_{Y_n} = \max \left\{ \frac{1}{n} \|y\|_{\ell_\infty}, \|Qy\|_{\ell_\infty/c_0} \right\}.$$

Note that Q remains a quotient map for the usual norm on ℓ_∞/c_0 .

Let $Z = (\sum Y_n)_{c_0}$, and assume that there is a uniformly continuous retraction of $B_{Z^{**}}$ onto B_Z . Then it follows that there is a sequence of retractions $g_n : Y^{**} \rightarrow Y_n$ which is equi-uniformly continuous, i.e., their moduli of uniform continuity satisfy

$$\psi_{g_n}(t) \leq \psi(t) \quad 0 < t \leq 2, \quad \text{with} \quad \lim_{t \rightarrow 0} \psi(t) = 0.$$

Consider the map $h_n : D_n \rightarrow Y_n$. If $x \neq x' \in D_n$, then

$$\|h_n(x) - h_n(x')\| \leq \max \left\{ \frac{2}{n}, \|x - x'\| \right\} \leq 2\|x - x'\|.$$

Since $B_{Y_n^{**}}$ is a 1-absolute Lipschitz retract, there is an extension $f_n : B_{\ell_\infty/c_0} \rightarrow B_{Y_n^{**}}$ of h_n with $\text{Lip}(f_n) \leq 2$. Now, if $x \in B_{\ell_\infty/c_0}$, there is $x' \in D_n$ with $\|x - x'\| < 2/n$. Thus,

$$\|g_n(f_n(x)) - g_n(f_n(x'))\| \leq \psi \left(\frac{4}{n} \right),$$

and hence,

$$\|Q(g_n(f_n(x))) - x\| \leq \psi \left(\frac{4}{n} \right) + \frac{2}{n}.$$

Then, for n large enough, we have

$$\psi \left(\frac{4}{n} \right) + \frac{2}{n} < 1.$$

By Lemma 5.7, this means that there is a uniformly continuous selection of the quotient map $Q : B_{\ell_\infty/c_0} \rightarrow Y_n$. Thus, B_{ℓ_∞/c_0} uniformly embeds into Y_n which is isomorphic to ℓ_∞ , and this is a contradiction with Theorem 5.2. \square

Problem 10. Does there exist a separable, or at least a weakly compactly generated (WCG) Banach space X which is not a Lipschitz retract of its bidual?

We note that, if X is Lipschitz embedded in ℓ_∞ , then X admits a countable separating family of Lipschitz real valued functions on X . Let us also recall that Bourgain proved in [16] that ℓ_∞/c_0 has no equivalent strictly convex norm. This somehow suggests the following problem.

Problem 11. If X is Lipschitz embeddable into ℓ_∞ , does X have an equivalent strictly convex norm?

We will finish this section by discussing a few more examples of non-isomorphic nonseparable spaces that have the same Lipschitz structure. We will discuss one way of getting many examples by using the so called

pull-back construction in the theory of exact sequences (see [21]). First, we need some preparation.

A diagram $0 \rightarrow Y \rightarrow X \rightarrow Z \rightarrow 0$ of Banach spaces and operators is said to be an exact sequence if the kernel of each arrow coincides with the image of the preceding one. Hence, in the diagram above, the second arrow denotes an injection and the third one is a quotient map. This means, by the open mapping theorem, that Y is isomorphic to a closed subspace of X and that the corresponding quotient is isomorphic to Z . We say that X is a twisted sum of Y and Z or an extension of Y by Z .

We say that the exact sequence *splits* if the second arrow i admits a linear retraction (i.e., an arrow r from X into Y so that $ri = \text{Id}_Y$) or, equivalently, if the third arrow q admits a linear section, or selection, i.e., if there is an arrow s from Z into X such that $qs = \text{Id}_Z$. This implies that then X is isomorphic to the direct sum $Y \oplus Z$.

Let $A : U \rightarrow Z$ and $B : V \rightarrow Z$ be two operators. The *pull-back* of $\{A, B\}$ is the space $PB = \{(u, v) : Au = Bv\} \subset U \times V$ considered with the canonical projections of $U \times V$ onto U and V , respectively.

If $0 \rightarrow Y \rightarrow X \rightarrow Z \rightarrow 0$ is an exact sequence with quotient map q , $T : V \rightarrow Z$ is an operator and PB denotes the pull-back of the couple $\{q, T\}$, then the diagram

$$\begin{array}{ccccccccc}
 0 & \longrightarrow & Y & \longrightarrow & X & \longrightarrow & Z & \longrightarrow & 0 \\
 & & & & \uparrow & & \uparrow & & \\
 0 & \longrightarrow & Y & \longrightarrow & PB & \longrightarrow & V & \longrightarrow & 0
 \end{array}$$

is commutative with exact rows. It follows that the pull-back sequence splits if and only if the operator T can be lifted to X , i.e., there exists an operator $\tau : V \rightarrow X$ such that $q\tau = T$.

Kalton's lemma [58] says that if the quotient map in the first row admits a Lipschitz section then so does the quotient map in the second row.

We now consider the following pull-back diagram.

$$\begin{array}{ccccccc}
 0 & \longrightarrow & c_0 & \longrightarrow & JL_\infty & \longrightarrow & c_0(I) \longrightarrow 0 \\
 & & \uparrow I_{c_0} & & \uparrow \hat{T} & & \uparrow T \\
 0 & \longrightarrow & c_0 & \longrightarrow & JL_2 & \longrightarrow & \ell_2(I) \longrightarrow 0 \\
 & & \uparrow I_{c_0} & & \uparrow \hat{S} & & \uparrow S \\
 0 & \longrightarrow & c_0 & \longrightarrow & CC & \longrightarrow & \ell_\infty \longrightarrow 0
 \end{array}$$

In this diagram, the operator T is the inclusion map and the operator S is the Rosenthal quotient map [97]. Therefore, we get that the space JL_2 of Johnson and Lindenstrauss is an example of a non WCG space, that it is Lipschitz isomorphic to $\ell_2(I) \oplus c_0$ and Lipschitz embeds into ℓ_∞ . However, it is known that it does not linearly isomorphically embed into ℓ_∞ (see [49]).

The space CC in the third row is then an example of a space that is Lipschitz homeomorphic to $\ell_\infty \oplus c_0$ but is not linearly isomorphic to it. Note also that CC is linearly isomorphic to a subspace of ℓ_∞ because JL_∞ linearly embeds into ℓ_∞ and CC is a subspace of $JL_\infty \oplus \ell_\infty$ (see [20] for details).

Similarly, Kalton obtained, for instance, that any nonseparable WCG space that contains an isomorphic copy of c_0 fails to have unique Lipschitz structure.

Problem 12. Does every reflexive (superreflexive) space have unique Lipschitz structure?

Problem 13. Does ℓ_∞ have unique Lipschitz structure?

Remark. It turns out that the undecidability of the continuum hypothesis (and of related statements) casts a shadow on the Lipschitz classification of non-separable spaces. Let I be a set of cardinality c . We assume that the space $c_0(I)$ is Lipschitz-isomorphic to a WCG Banach space, X . Does it follow that X is linearly isomorphic to $c_0(I)$? It follows from [38] that the answer is positive if $c < \aleph_{\omega_0}$, where \aleph_{ω_0} denotes the ω_0 th cardinal. On the other hand, it follows from [11, 73] that the answer is negative if $c > \aleph_{\omega_0}$ (see also [7] for a related result on the failure of a non-separable Sobczyk theorem at the density \aleph_{ω_0}).

Since (ZFC) does not decide if c is strictly less or strictly more than \aleph_{ω_0} , the above question is undecidable in (ZFC).

6. Coarse embeddings into Banach spaces and geometric group theory. Two topological spaces M and N are homotopically equivalent if there exist continuous maps $f : M \rightarrow N$ and $g : N \rightarrow M$ such that $f \circ g$ and $g \circ f$ are both homotopic to the identity map on the space on which they operate. For instance, any topological *vector* space is homotopically equivalent to a point (in other words, is contractible).

We consider the case where M and N are real compact manifolds. In order to understand their geometry, it is, of course, important to find quantities which are invariant under homotopy equivalence. A basic example is the signature of the manifold, which can be defined as follows when the dimension $n = 4k$ is divisible by 4. If d_j denotes the exterior derivative acting on the differential forms of degree j , then $F_k = \text{Ker}(d_{2k})$ is the vector space of closed forms of degree $2k$ and $E_k = \text{Im}(d_{2k-1})$ is its subspace of exact forms of degree $2k$. If ω_1 and ω_2 are two forms in F_k and, if we define Q by

$$Q(\omega_1, \omega_2) = \int_M \omega_1 \wedge \omega_2,$$

then Q is a bilinear symmetric form, and an easy computation shows that the value of Q depends only upon the classes of ω_1 and ω_2 in the finite-dimensional quotient space $H_k(M) = F_k/E_k$. Therefore, Q defines a quadratic form on $H_k(M)$, and its signature is called the signature of the manifold M . This signature is invariant under homotopy equivalence. Actually, a theorem of Novikov asserts (for simply connected manifolds) that the signature is the only homotopy invariant which can be computed in terms of quantities called Pontryagin polynomials.

We now recall the basics of geometric group theory. Let G be a finitely generated group and S a finite set generating G . We can equip G with the word distance d_S associated to S , as follows: if $\|g\|_S$ is the minimal length of a word written with elements of S and S^{-1} representing g , then we define the left-invariant distance d_S on G by $d_S(g_1, g_2) = \|g_1^{-1}g_2\|_S$. If, for instance, G is the group \mathbb{Z}^n and S is its generating set consisting of the unit vector basis, then d_S coincides with the distance induced by the ℓ_1 norm on \mathbb{R}^n . It is easily checked that if S

and S' are two finite generating sets, then the identity map is a coarse Lipschitz isomorphism between the metric spaces (G, d_S) and $(G, d_{S'})$. A property of finitely generated groups is said to be *geometric* if it depends only upon the space (G, d_S) up to coarse Lipschitz equivalence. Many natural properties of groups (such as amenability, hyperbolicity, or being virtually nilpotent, i.e., containing a nilpotent subgroup of finite index) turn out to be geometric.

We denote again by M a real compact manifold. Novikov's conjecture asserts that certain "higher signatures" are homotopy invariants. We would have to introduce several highly non-trivial concepts before providing a precise statement of this conjecture, and this is not the purpose of this survey. However, it is easy to describe the link between Novikov's conjecture and geometric group theory. Indeed, let $\pi_1(M)$ be the first homotopy group of the manifold M . Gromov conjectured a link between the geometry of the finitely generated group $\pi_1(M)$ and the Novikov conjecture, and this conjecture was confirmed by Yu: Novikov's conjecture (and even the stronger coarse Baum-Connes conjecture) holds true if the group $\pi_1(M)$ equipped with the word distance coarsely embeds into the Hilbert space [100]. This important result was later generalized by Kasparov and Yu [64] who showed that the Hilbert space can be replaced in this statement by any super-reflexive space. This is indeed a generalization since the spaces ℓ_p with $p > 2$ do not coarsely embed into the Hilbert space [53]. Note that, conversely, it is an open question to know if the separable Hilbert space coarsely embeds into every infinite-dimensional Banach space; in other words, if coarse embedding into ℓ_2 is the strongest possible property of that kind. A word of warning is needed here: what we call "coarse embedding" is often called (after Gromov) "uniform embedding" in the context of differential geometry. In this survey, however, the word uniform bears another meaning.

Let us recall that a metric space E is called locally finite if every ball of E is finite, and it has *bounded geometry* if, for any $r > 0$, the cardinality of subsets of diameter less than r is uniformly bounded. The left invariance of the distance d_S shows that any finitely generated group has bounded geometry. Therefore, the question occurs of deciding which finitely generated group, and more generally which space with bounded geometry coarsely embeds into a super-reflexive space. For instance, could it be that every space with bounded

geometry coarsely embeds into a super-reflexive space?

This question is now negatively answered. A first example of a locally finite space which does not coarsely embed into the Hilbert space is obtained in [25], using in particular a construction of Enflo [30]. Then Gromov shows [42], through a random approach, the existence of finitely generated groups G such that the metric space (G, d_G) “somewhat coarsely” (see [46, pages 350–351]) contains a sequence of expanders (E_i) such that the girth of E_i , namely, the length of the shortest closed curve, increases to infinity. As shown in [72], embeddings of expanders into the Hilbert space have maximal distortion. It follows that G cannot be coarsely embedded into a Hilbert space. Since such a group can be realized as a homotopy group (see [43]), Gromov’s result shows that attacking the full Novikov conjecture through the coarse embedding approach ought to be difficult (see also [46]). Later, Lafforgue obtained by an algebraic approach a sequence of expanders which does not coarsely embed into a super-reflexive space [66], while Mendel and Naor obtained such expanders by a completely different method [79]. The existence of spaces with bounded geometry which do not coarsely embed into a super-reflexive space follows. Let us mention that Lafforgue showed that his sequence of expanders does not coarsely embed into a space with non trivial type ([67]; see also [69] and [70] for recent progress along these lines).

On the other hand, the problem remains open to decide which metric spaces coarsely embed into Banach spaces of given regularity. Any metric space with bounded geometry coarsely embeds into a reflexive space [19]. This result is widely generalized in [56], where it is shown that every stable metric space (where “stable” means that the order of limits can be permuted in $\lim_k \lim_n d(x_k, y_n)$ each time all limits exist) can be coarsely embedded into a reflexive space. This is indeed extending [19] since every metric space whose balls are compact, and thus every locally finite metric space, is stable. Moreover, it follows from Theorem 4.4, that any locally finite metric space Lipschitz embeds into the following very simple reflexive space: $(\sum_{n=1}^{\infty} \ell_{\infty}^n)_{\ell_2}$ (note that this space is both AUC and AUS). On the other hand, Theorem 4.7 states that c_0 does not coarsely embed into a reflexive space, nor into a stable metric space (by the above, or directly by [94]). Note that the important stable Banach space L^1 coarsely embeds into the Hilbert space. More precisely, Schoenberg proved in [99] that $(L^1, \|\cdot\|_1^{1/2})$ is

isometric to a subset of L^2 .

As seen before, coarse embeddings of special graphs bear important consequences on the non-linear geometry of Banach spaces. Sequences of expanders shed light on the geometry of groups and its applications to homotopy invariants. It is plausible that such expanders could provide several interesting examples in geometry of Banach spaces.

For the record, we recall the following:

Problem 14. Does ℓ_2 coarsely embed into every infinite dimensional Banach space?

Note that the answer to this problem is positive for spaces with an unconditional basis and non trivial cotype (see [83, 87]). The proof is based on the work of Odell and Schlumprecht on sphere homeomorphisms [85] and on a criterion of Dadarlat and Guentner for coarse embeddability into a Hilbert space [23].

7. Lipschitz-free spaces and their applications. Let M be a pointed metric space, that is, a metric space equipped with a distinguished point denoted 0 . The space $\text{Lip}_0(M)$ is the space of real-valued Lipschitz functions on M which vanish at 0 . Let $\mathcal{F}(M)$ be the natural predual of $\text{Lip}_0(M)$, whose w^* -topology coincides on the unit ball of $\text{Lip}_0(M)$ with the pointwise convergence on M . The Dirac map $\delta : M \rightarrow \mathcal{F}(M)$ defined by $\langle g, \delta(x) \rangle = g(x)$ is an isometric embedding from M to a subset of $\mathcal{F}(M)$ which generates a dense linear subspace. This predual $\mathcal{F}(M)$ is called in [37] the *Lipschitz-free space over M* . When M is separable, $\mathcal{F}(M)$ is separable as well since $\delta(M)$ spans a dense subspace. Although Lipschitz-free spaces over separable metric spaces constitute a class of separable Banach spaces which are easy to define, the structure of these spaces is very poorly understood to this day. Improving our understanding of this class is a fascinating research program. Note that, if we identify (through the Dirac map) a metric space M with a subset of $\mathcal{F}(M)$, any Lipschitz map from M to a metric space N extends to a continuous linear map from $\mathcal{F}(M)$ to $\mathcal{F}(N)$. So Lipschitz maps become linear, but of course the complexity is shifted from the map to the free space: this may explain why the structure of Lipschitz-free spaces is not easy to analyze. A first example is provided by the real line, whose free space is isometric to L_1 . Actually, metric

spaces M whose free space is isometric to a subspace of L_1 are characterized in [34] as subsets of metric trees equipped with the least path metric. On the other hand, the free space of the plane \mathbb{R}^2 does not embed isomorphically into L^1 [82]. Note that the Lipschitz free spaces are often called Arens-Eells spaces and that they can be interpreted in terms of “transportation cost.”

Banach spaces X are, in particular, pointed metric spaces (pick the origin as distinguished point) and we can apply the previous construction. Note that the isometric embedding $\delta : X \rightarrow \mathcal{F}(X)$ is, of course, non linear since there exist Lipschitz functions on X which are not affine.

This Dirac map has a linear left inverse $\beta : \mathcal{F}(X) \rightarrow X$ which is the quotient map such that $x^*(\beta(\mu)) = \langle x^*, \mu \rangle$ for all $x^* \in X^*$; in other words, β is the extension to $\mathcal{F}(X)$ of the barycenter map. This setting provides canonical examples of Lipschitz-isomorphic spaces. Indeed, if we let $Z_X = \text{Ker}(\beta)$, it follows easily from $\beta\delta = \text{Id}_X$ that $Z_X \oplus X = \mathcal{G}(X)$ is Lipschitz-isomorphic to $\mathcal{F}(X)$.

Following [37], let us say that a Banach space X has the *lifting property* if there is a continuous linear map $R : X \rightarrow \mathcal{F}(X)$ such that $\beta R = \text{Id}_X$, or equivalently, if for Y and Z Banach spaces and $S : Z \rightarrow Y$ and $T : X \rightarrow Y$ continuous linear maps, the existence of a Lipschitz map \mathcal{L} such that $T = S\mathcal{L}$ implies the existence of a continuous linear operator L such that $T = SL$. A diagram-chasing argument shows that $\mathcal{G}(X)$ is linearly isomorphic to $\mathcal{F}(X)$ if and only if X has the lifting property [37]. It turns out that all non-separable reflexive spaces, and also the spaces $\ell_\infty(\mathbb{N})$ and $c_0(\Gamma)$ when Γ is uncountable, fail the lifting property and this provides canonical examples of pairs of Lipschitz but not linearly isomorphic spaces.

On the other hand, the following result is proved in [37].

Theorem 7.1. *Every separable Banach space X has the lifting property.*

Proof. We will actually give two proofs. First, one can pick a Gaussian measure γ whose support is dense in X and use the result that $(\delta * \gamma)$ is Gâteaux-differentiable; then, in the above notation, $R = (\delta * \gamma)'(0)$ satisfies $\text{Id}_X = \beta R$.

The second proof is essentially self-contained. It consists of replacing the Gaussian measure by a cube measure, and this will be useful later. It underlines the simple fact that being separable is equivalent to being “compact-generated.” Again, we use differentiation, but only in the directions which are normal to the faces of the cube.

Let $(x_i)_{i \geq 1}$ be a linearly independent sequence of vectors in X such that

$$\overline{\text{span}}[(x_i)_{i \geq 1}] = X$$

and $\|x_i\| = 2^{-i}$ for all i . Let $H = [0, 1]^{\mathbb{N}}$ be the Hilbert cube and $H_n = [0, 1]^{\mathbb{N}_n}$ the copy of the Hilbert cube where the factor of rank n is omitted; that is, $\mathbb{N}_n = \mathbb{N} \setminus \{n\}$. We denote by λ (respectively, λ_n) the natural probability measure on H (respectively, H_n) obtained by taking the product of the Lebesgue measure on each factor.

We denote $E = \text{span}[(x_i)_{i \geq 1}]$ and $R : E \rightarrow \mathcal{F}(X)$ the unique linear map which satisfies for all $n \geq 1$ et all $f \in \text{Lip}_0(X)$

$$R(x_n)(f) = \int_{H_n} \left[f \left(x_n + \sum_{j=1, j \neq n}^{\infty} t_j x_j \right) - f \left(\sum_{j=1, j \neq n}^{\infty} t_j x_j \right) \right] d\lambda_n(t).$$

Pick $f \in \text{Lip}_0(X)$. If the function f is Gâteaux-differentiable, Fubini’s theorem shows that, for all $x \in E$,

$$R(x)(f) = \int_H \langle \{\nabla f\} \left(\sum_{j=1}^{\infty} t_j x_j \right), x \rangle d\lambda(t).$$

Thus, $|R(x)(f)| \leq \|x\| \|f\|_L$ in this case. But, since X is separable, any $f \in \text{Lip}_0(X)$ is a uniform limit of a sequence (f_j) of Gâteaux-differentiable functions such that $\|f_j\|_L \leq \|f\|_L$. It follows that

$$\|R\| \leq 1.$$

We may now extend R to a linear map $\overline{R} : X \rightarrow \mathcal{F}(X)$ such that $\|\overline{R}\| = 1$, and it is clear that $\overline{R}(x)(x^*) = x^*(x)$ for all $x \in X$ and all $x^* \in X^*$. Hence, $\beta \overline{R} = \text{Id}_X$. \square

The above proof follows [37]. We refer to [35] for an elementary approach along the same lines, which uses only finite-dimensional arguments and is accessible at the undergraduate level.

The lifting property for separable spaces forbids the existence of a *separable* Banach space X such that $\mathcal{F}(X)$ and $\mathcal{G}(X)$ are not linearly isomorphic, but on the other hand it shows that if there exists an isometric embedding from a separable Banach space X into a Banach space Y , then Y contains a linear subspace which is isometric to X . Indeed, a theorem due to Figiel [33] states that, if $J : X \rightarrow Y$ is an isometric embedding such that $J(0) = 0$ and $\overline{\text{span}}[J(X)] = Y$, then there is a linear quotient map with $\|Q\| = 1$ and $QJ = \text{Id}_X$, the lifting property provides a linear contractive map R such that $QR = \text{Id}_X$, and this map R is a linear isometric embedding. We note that $P = RQ$ is a contractive projection from Y onto $R(X)$. This remark is developed further in [36], where it is shown that the existence of a non-linear isometric embedding from X into Y is a very restrictive condition on the couple (X, Y) .

Nigel Kalton constructed the proper frame for showing the gap which separates Hölder maps from Lipschitz ones [55]. If $(X, \|\cdot\|)$ is a Banach space and $\omega : [0, +\infty) \rightarrow [0, +\infty)$ is a subadditive function such that $\lim_{t \rightarrow 0} \omega(t) = \omega(0) = 0$ and $\omega(t) = t$ if $t \geq 1$, then the space $\text{Lip}_\omega(X)$ of $(\omega \circ d)$ -Lipschitz functions on X which vanish at 0 has a natural predual denoted $\mathcal{F}_\omega(X)$, and the barycentric map $\beta_\omega : \mathcal{F}_\omega(X) \rightarrow X$ (whose adjoint is the canonical embedding from X^* to $\mathcal{F}_\omega(X)$) is still a linear quotient map such that $\beta_\omega \delta = \text{Id}_X$. However, the Dirac map $\delta : X \rightarrow \mathcal{F}_\omega(X)$ is now uniformly continuous with modulus ω , e.g., α -Hölder when $\omega(t) = \max(t^\alpha, t)$ with $0 < \alpha < 1$. Uniformly continuous functions fail the differentiability properties that Lipschitz functions enjoy, and thus one can expect that this part of the theory is more “distant” from the linear theory than the Lipschitz one. It is indeed so, and [55, Theorem 4.6], reads as follows.

Theorem 7.2. *If ω satisfies $\lim_{t \rightarrow 0} \omega(t)/t = \infty$, then $\mathcal{F}_\omega(X)$ is a Schur space, that is, weakly convergent sequences in $\mathcal{F}_\omega(X)$ are norm convergent.*

It follows from Theorem 7.2 that the uniform analogue of the lifting property fails unless X has the (quite restrictive) Schur property. Moreover, $\mathcal{F}_\omega(X)$ is (3ω) -uniformly homeomorphic to $[X \oplus \text{Ker}(\beta_\omega)]$ and as soon as $\lim_{t \rightarrow 0} \omega(t)/t = 0$ and X fails the Schur property, we obtain canonical pairs of uniformly (even Hölder) homeomorphic

separable Banach spaces which are not linearly isomorphic. We refer to [51, 95] for other examples of such pairs.

Along with Hölder maps between Banach spaces, one may as well consider Lipschitz maps between quasi-Banach spaces, and this is done in [6] where similar methods provide examples of separable quasi-Banach spaces which are Lipschitz but not linearly isomorphic.

We now observe that the proof (with cube measures) of Theorem 7.1 provides the existence of compact metric spaces whose free space fails the approximation property (in short, AP). This has been observed in [40].

Theorem 7.3. *There exists a compact metric space K whose free space $\mathcal{F}(K)$ fails the approximation property.*

Proof. We use the notation of the proof of Theorem 7.1. Let C be the closed convex hull of the sequence $(x_i)_{i \geq 1}$, and let $K = 2C$. It is easily seen that the map R takes its values in the closed subspace $\mathcal{F}(K)$ of $\mathcal{F}(X)$, and so does \bar{R} . It follows that X is isometric to a 1-complemented subspace of $\mathcal{F}(K)$, through the projection $\bar{R}Q$. If this construction is applied to a Banach space X which fails AP, then $\mathcal{F}(K)$ fails AP as well since AP is carried to complemented subspaces. \square

Problem 15. Let X be a separable Banach space and Y a Banach space which is Lipschitz-isomorphic to X . Does it follow that Y is linearly isomorphic to X ?

This question amounts to knowing whether every separable Banach space is determined by its metric structure. It is open, for instance, if $X = \ell_1$ or if $X = C(K)$ with K a countable compact metric space, unless $C(K)$ is isomorphic to c_0 . Note that, by the above, the answer to this question is negative if we drop the separability assumption, if we replace Lipschitz by Hölder, or if we replace Banach by quasi-Banach.

Problem 16. Is the Lipschitz-free space $\mathcal{F}(\ell_1)$ over ℓ_1 complemented in its bidual?

A motivation for this question is that, if $\mathcal{F}(\ell_1)$ is complemented in its bidual, it follows that every space X which is Lipschitz-isomorphic

to ℓ_1 is complemented in its bidual, and then [13, Corollary 7.7] shows that X is linearly isomorphic to ℓ_1 .

Problem 17. Theorem 7.3 leads to the question of knowing for which compact spaces K the space $\mathcal{F}(K)$ has AP or its metric version MAP. So far, very little is known on this topic, which is related with the existence of linear extension operators for Lipschitz functions (see [15, 40, 68]).

Problem 18. Let M be an arbitrary uniformly discrete metric space, that is, there exists $\theta > 0$ such that $d(x, y) \geq \theta$ for all $x \neq y$ in M . Does $\mathcal{F}(M)$ have the BAP? Note that AP holds by ([55, Proposition 4.4]). A positive answer to this question would imply that every separable Banach space X is approximable, that is, the identity Id_X is the pointwise limit of an equi-uniformly continuous sequence of maps with relatively compact range. By [59, Theorem 4.6], it is indeed so for X and X^* when X^* is separable, and in particular, every separable reflexive space is approximable. On the other hand, a negative answer to this question would provide an equivalent norm on ℓ_1 failing MAP, and this would solve a famous problem in approximation theory, by providing the first example of a dual space—namely, ℓ_∞ equipped with the corresponding dual norm—with AP (and even BAP) but failing MAP.

Acknowledgments. The authors would like to thank F. Baudier, M. Kraus and V. Montesinos for useful discussions. They also wish to thank M.I. Ostrovskii for his valuable suggestions that helped to improve this survey.

REFERENCES

1. I. Aharoni, *Every separable metric space is Lipschitz equivalent to a subset of c_0^+* , Israel J. Math. **19** (1974), 284–291.
2. I. Aharoni and J. Lindenstrauss, *Uniform equivalence between Banach spaces*, Bull. Amer. Math. Soc. **84** (1978), 281–283.
3. F. Albiac, *Non-linear structure of some classical quasi-Banach spaces and F -spaces*, J. Math. Anal. Appl. **340** (2008), 1312–1325.
4. F. Albiac and F. Baudier, *Embeddability of snowflaked metrics with applications to the nonlinear geometry of the spaces L_p and ℓ_p for $0 < p < \infty$* , J. Geom. Anal., to appear, doi:10.1007/s12220-013-9390-0.

5. F. Albiac and N.J. Kalton, *Topics in Banach space theory*, Grad. Texts Math. **233**, Springer, New York, 2006.
6. ———, *Lipschitz structure of quasi-Banach spaces*, Israel J. Math. **170** (2009), 317–335.
7. S. Argyros, J. Castillo, A. Granero, M. Jimenez and J. Moreno, *Complementation and embeddings of $c_0(I)$ in Banach spaces*, Proc. Lond. Math. Soc. **85** (2002), 742–768.
8. F. Baudier, *Metricl characterization of super-reflexivity and linear type of Banach spaces*, Arch. Math. **89** (2007), 419–429.
9. F. Baudier, N.J. Kalton, G. Lancien, *A new metric invariant for Banach spaces*, Stud. Math. **199** (2010), 73–94.
10. F. Baudier and G. Lancien, *Embeddings of locally finite metric spaces into Banach spaces*, Proc. Amer. Math. Soc. **136** (2008), 1029–1033.
11. M. Bell and W. Marciszewski, *On scattered Eberlein compact spaces*, Israel J. Math. **158** (2007), 217–224.
12. Y. Benyamini, *An M -space which is not isomorphic to a $C(K)$ space*, Israel J. Math. **28** (1977), 98–102.
13. Y. Benyamini and J. Lindenstrauss, *Geometric nonlinear functional analysis*, A.M.S. Colloq. Publ. **48**, American Mathematical Society, Providence, RI, 2000.
14. L. Borel-Mathurin, *The Szlenk index of Orlicz sequence spaces*, Proc. Amer. Math. Soc. **138** (2010), 2043–2050.
15. ———, *Approximation properties and non-linear geometry of Banach spaces*, Houston J. Math. **38** (2012), 1135–1148.
16. J. Bourgain, *ℓ_∞/c_0 has no strictly convex norm*, Proc. Amer. Math. Soc. **78** (1985), 225–226.
17. ———, *The metricl interpretation of superreflexivity in Banach spaces*, Israel J. Math. **56** (1986), 222–230.
18. J. Bourgain, V. Milman and H. Wolfson, *On type of metric spaces*, Trans. Amer. Math. Soc. **294** (1986), 295–317.
19. N. Brown and E. Guentner, *Uniform embeddings of bounded geometry spaces into reflexive Banach space*, Proc. Amer. Math. Soc. **133** (2005), 2045–2050.
20. F. Cabello Sánchez and J.M.F. Castillo, *Uniform boundedness and twisted sums of Banach spaces*, Houst. J. Math. **30** (2004), 523–536.
21. F. Cabello Sánchez, J.M.F. Castillo, N.J. Kalton and D.T. Yost, *Twisted sums with $C(K)$ -spaces*, Trans. Amer. Math. Soc. **355** (2003), 4523–4541.
22. P.G. Casazza, *James' quasi-reflexive space is primary*, Israel J. Math. **26** (1977), 294–305.
23. M. Dadarlat and E. Guentner, *Constructions preserving Hilbert space uniform embeddability of discrete groups*, Trans. Amer. Math. Soc. **355** (2003), 3253–3275.
24. R. Deville, G. Godefroy and V. Zizler, *Smoothness and renormings in Banach spaces*, Pitman Mono. Surv. **64**, Longman, New York, 1993.

25. A.N. Dranishnikov, G. Gong, V. Lafforgue and G. Yu, *Uniform embeddings into Hilbert space and a question of Gromov*, Canad. Math. Bull. **45** (2002), 60–70.
26. Y. Dutrioux, *Quotients of c_0 and Lipschitz-homeomorphisms*, Houst. J. Math. **27** (2001), 593–599.
27. ———, *Géométrie non linéaire des espaces de Banach*, Ph.D. Thesis, l'Université Paris VI, 2002.
28. Y. Dutrioux and G. Lancien, *Isometric embeddings of compact spaces into Banach spaces*, J. Funct. Anal. **255** (2008), 494–501.
29. I.S. Èdelštein and P. Wojtaszczyk, *On projections and unconditional bases in direct sums of Banach spaces*, Stud. Math. **56** (1976), 263–276.
30. P. Enflo, *On a problem of Smirnov*, Ark. Mat. **8** (1969), 107–109.
31. ———, *On infinite-dimensional topological groups*, Ecole Polytech., Palaiseau, 1978.
32. M. Fabian, P. Habala, P. Hájek, V. Montesinos and V. Zizler, *Banach space theory. The basis for linear and nonlinear analysis*, CMS Books Math., Springer, New York, 2011.
33. T. Figiel, *On non linear isometric embeddings of normed linear spaces*, Bull. Acad. Polon. Sci. Math. Astro. Phys. **16** (1968), 185–188.
34. A. Godard, *Tree metrics and their Lipschitz-free spaces*, Proc. Amer. Math. Soc. **138** (2010), 4311–4320.
35. G. Godefroy, *Linearization of isometric embeddings between Banach spaces: An elementary approach*, Opera. Matr. **6** (2012), 339–345.
36. ———, *Remarks on non-linear embeddings between Banach spaces*, Houst. J. Math. **26** (2010), 283–287.
37. G. Godefroy and N.J. Kalton, *Lipschitz-free Banach spaces*, Stud. Math. **159** (2003), 121–141.
38. G. Godefroy, N.J. Kalton and G. Lancien, *Subspaces of $c_0(\mathbb{N})$ and Lipschitz isomorphisms*, Geom. Funct. Anal. **10** (2000), 798–820.
39. ———, *Szlenk indices and uniform homeomorphisms*, Trans. Amer. Math. Soc. **353** (2001), 3895–3918.
40. G. Godefroy and N. Ozawa, *Free Banach spaces and the approximation properties*, Proc. Amer. Math. Soc., to appear.
41. E. Gorelik, *The uniform nonequivalence of L_p and l_p* , Israel J. Math. **87** (1994), 1–8.
42. M. Gromov, *Random walks in random groups*, Geom. Funct. Anal. **13**(2003), 73–146.
43. ———, *Spaces and questions*, Geom. Funct. Anal., Special Volume-GAFA **2000**, 118–161.
44. R. Haydon, E. Odell and T. Schlumprecht, *Small subspaces of L_p* , Ann. Math. **173** (2011), 169–209.
45. S. Heinrich and P. Mankiewicz, *Applications of ultrapowers to the uniform and Lipschitz classification of Banach spaces*, Stud. Math. **73** (1982), 225–251.

46. N. Higson, V. Lafforgue and G. Skandalis, *Counterexamples to the Baum-Connes conjecture*, Geom. Funct. Anal. **12** (2002), 330–354.
47. R.C. James, *Uniformly non-square Banach spaces*, Ann. Math. **80** (1964), 542–550.
48. W.B. Johnson, *Operators into L_p which factor through ℓ_p* , J. Lond. Math. Soc. **14** (1976), 333–339.
49. W.B. Johnson and J. Lindenstrauss, *Some remarks on weakly compactly generated Banach spaces*, Israel J. Math. **17**, (1974), 219–230.
50. W. B. Johnson, J. Lindenstrauss, D. Preiss and G. Schechtman, *Almost Fréchet differentiability of Lipschitz mappings between infinite dimensional Banach spaces*, Proc. Lond. Math. Soc. **84** (2002), 711–746.
51. W.B. Johnson, J. Lindenstrauss and G. Schechtman, *Banach spaces determined by their uniform structures*, Geom. Funct. Anal. **6** (1996), 430–470.
52. W.B Johnson and E. Odell, *Subspaces of L_p which embed into ℓ_p* , Compos. Math. **28** (1974), 37–49.
53. W.B. Johnson and N.L. Randrianarivony, *l_p ($p > 2$) does not coarsely embed into a Hilbert space*, Proc. Amer. Math. Soc. **134** (2006), 1045–1050.
54. W.B Johnson and G. Schechtman, *Diamond graphs and super-reflexivity*, J. Topol. Anal. **1** (2009), 177–189.
55. N.J. Kalton, *Spaces of Lipschitz and Hölder functions and their applications*, Collect. Math. **55** (2004), 171–217.
56. ———, *Coarse and uniform embeddings into reflexive spaces*, Quart. J. Math. (Oxford) **58** (2007), 393–414.
57. ———, *The nonlinear geometry of Banach spaces*, Rev. Mat. Comp. **21**, (2008), 7–60.
58. ———, *Lipschitz and uniform embeddings into ℓ_∞* , Fund. Math. **212** (2011), 53–69.
59. ———, *The uniform structure of Banach spaces*, Math. Ann. **354** (2012), 1247–1288.
60. ———, *Uniform homeomorphisms of Banach spaces and asymptotic structure*, Trans. Amer. Math. Soc. **365** (2013), 1051–1079.
61. ———, *Examples of uniformly homeomorphic Banach spaces*, Israel J. Math., to appear.
62. N.J. Kalton and G. Lancien, *Best constants for Lipschitz embeddings of metric spaces into c_0* , Fund. Math. **199** (2008), 249–272.
63. N.J. Kalton and N.L. Randrianarivony, *The coarse Lipschitz structure of $\ell_p \oplus \ell_q$* , Math. Ann. **341** (2008), 223–237.
64. G. Kasparov and G. Yu, *The coarse geometric Novikov conjecture and uniform convexity*, Adv. Math. **206** (2006), 1–56.
65. M. Kraus, *Coarse and uniform embeddings between Orlicz sequence spaces*, Mediterr. J. Math., to appear.

66. V. Lafforgue, *Un renforcement de la propriété (T)*, Duke Math. J. **143** (2008), 559–602.
67. ———, *Propriété (T) renforcée banachique et transformée de Fourier rapide*, J. Topol. Anal. **1** (2009), 191–206.
68. G. Lancien and E. Pernecká, *Approximation properties and Schauder decompositions in Lipschitz-free spaces*, J. Funct. Anal., to appear.
69. B. Liao, *Strong Banach property (T), after Vincent Lafforgue*, arXiv: 1212.4646 (2012).
70. ———, *Strong Banach property (T) for simple algebraic groups of higher rank*, arXiv: 1301.1861 (2013).
71. J. Lindenstrauss, *On nonlinear projections in Banach spaces*, Michigan Math. J. **11** (1964), 263–287.
72. N. Linial, E. London and Y. Rabinovich, *The geometry of graphs and some of its algorithmic applications*, Combinatorica **15** (1995), 215–245.
73. W. Marciszewski, *On Banach spaces $C(K)$ isomorphic to $c_0(\Gamma)$* , Stud. Math. **156** (2003), 295–302.
74. B. Maurey and G. Pisier, *Séries de variables aléatoires vectorielles indépendantes et propriétés géométriques des espaces de Banach*, Stud. Math. **58** (1976), 45–90.
75. S. Mazur and S. Ulam, *Sur les transformations isométriques d’espaces vectoriels normés*, C.R.A.S. Paris, **194**, (1932), 946–948.
76. M. Mendel and A. Naor, *Euclidean quotients of finite metric spaces*, Adv. Math. **189** (2004), 451–494.
77. ———, *Scaled Enflo type is equivalent to Rademacher type*, Bull. Lond. Math. Soc. **39** (2007), 493–498.
78. ———, *Metric cotype*, Ann. Math. **168** (2008), 247–298.
79. ———, *Nonlinear spectral calculus and super-expanders*, preprint (2012), arXiv:1207.4705, Publ. Math. IHÉS, to appear. Conference version and extended abstract in: *Towards a calculus for non-linear spectral gaps*, SODA 10: Proc. 21st annual ACM-SIAM Symposium on Discrete Algorithms, M. Charikar, ed., SIAM, 2010.
80. V.D. Milman, *Geometric theory of Banach spaces, II, Geometry of the unit ball*, Uspehi Mat. Nauk **26** (1971), 73–149. Russian Math. Surv. **26** (1971), 79–163 (in English).
81. A. Naor, *An introduction to the Ribe program*, Japanese J. Math. **7** (2012), 167–233.
82. A. Naor and G. Schechtman, *Planar earthmover is not in L^1* , SIAM J. Comp. **37** (2007), 804–826.
83. P. Nowak, *On coarse embeddability into ℓ_p -spaces and a conjecture of Dranishnikov*, Fund. Math. **189** (2006), 111–116.
84. ———, *Coarse embeddings of metric spaces into Banach spaces*, Proc. Amer. Math. Soc. **133** (2005), 2589–2596.

- 85.** E. Odell and T. Schlumprecht, *The distortion problem*, Acta Math. **173** (1994), 259–281.
- 86.** ———, *Embeddings into Banach spaces with finite dimensional decompositions*, Rev. Real Acad. Cienc. **100** (2006), 295–323.
- 87.** M.I. Ostrovskii, *Coarse embeddability into Banach spaces*, Topol. Proc. **33** (2009), 163–183.
- 88.** ———, *Embeddability of locally finite metric spaces into Banach spaces is finitely determined*, Proc. Amer. Math. Soc. **140** (2012), 2721–2730.
- 89.** ———, *Different forms of metric characterizations of classes of Banach spaces*, Houst. J. Math., to appear.
- 90.** I.I. Parovičenko, *On a universal bicomactum of weight \aleph* , Dokl. Akad. Nauk SSSR **150** (1963), 36–39.
- 91.** A. Pełczyński, *Projections in certain Banach spaces*, Stud. Math. **19** (1960), 209–228.
- 92.** G. Pisier, *Martingales with values in uniformly convex spaces*, Israel J. Math. **20** (1975), 326–350.
- 93.** N.L. Randrianarivony, *Characterization of quasi-Banach spaces which coarsely embed into a Hilbert space*, Proc. Amer. Math. Soc. **134** (2006), 1315–1317.
- 94.** Y. Raynaud, *Espaces de Banach superstables, distances stables et homéomorphismes uniformes*, Israel J. Math. **44** (1983), 33–52.
- 95.** M. Ribe, *Existence of separable uniformly homeomorphic non isomorphic Banach spaces*, Israel J. Math. **48**, (1984), 139–147.
- 96.** ———, *On uniformly homeomorphic normed spaces*, Ark. Mat. **16** (1978), 1–9.
- 97.** H.P. Rosenthal, *On quasi-complemented subspaces of Banach spaces*, Proc. Natl. Acad. Sci. **59** (1968), 361–364.
- 98.** G. Schechtman, personal communication, **date?**_____.
- 99.** I.J. Schoenberg, *Metric spaces and positive definite functions*, Trans. Amer. Math. Soc. **44** (1938), 522–553.
- 100.** G. Yu, *The coarse Baum-Connes conjecture for spaces which admit a uniform embedding into Hilbert space*, Invent. Math. **139** (2000), 201–240.

INSTITUT DE MATHÉMATIQUES DE JUSSIEU, 4 PLACE JUSSIEU, 75005 PARIS, FRANCE
Email address: godefroy@math.jussieu.fr

UNIVERSITÉ DE FRANCHE-COMTÉ, LABORATOIRE DE MATHÉMATIQUES UMR 6623,
 16 ROUTE DE GRAY, 25030 BESANÇON CEDEX, FRANCE
Email address: gilles.lancien@univ-fcomte.fr

DEPARTMENT OF MATHEMATICAL SCIENCES, UNIVERSITY OF ALBERTA, CENTRAL
 ACADEMIC BUILDING, T6G 2G1 EDMONTON, ALBERTA, CANADA
Email address: vasekzizler@gmail.com